

Her setter du inn et bilde som illustrerer hva rapporten handler om.

Bildet må være i størrelsen 1250x836 piksler.

Slik setter du inn bilde:

1. Marker denne illustrasjonen
2. Klikk høyre musetast
3. Velg «Endre bilde»
4. Finn det bildet du vil sette inn



VKM Report 2021: XX

Assessment and updating of a fortification model

Scientific Opinion of the Panel on Nutrition, Dietetic products, Novel food and Allergy of the Norwegian Scientific Committee for Food and Environment

VKM Report 2021: XX
Assessment and updating of a fortification model

Scientific Opinion of the Panel on Nutrition, Dietetic products, Novel food and Allergy of the
Norwegian Scientific Committee for Food and Environment
XX.XX.2021

ISBN: 00000-00000
ISSN: 2535-4019
Norwegian Scientific Committee for Food and Environment (VKM)
Postboks 222 Skøyen
0213 Oslo
Norway

Phone: +47 21 62 28 00
Email: vkm@vkm.no

vkm.no

Cover photo: xx

Suggested citation: VKM, Lene Frost Andersen, Inger Therese Lillegaard, Bente Mangschou, Lisbeth Dahl, Knut Tomas Dalen, Kristin Holvik, Martinus Løvik, Christine L Parr, Tonje Holte Stea, Stine M Ulven, Tor A Strand (forfattere som oppfyller VKMs forfatterskapskriterier i følgende rekkefølge: faglig leder; medlemmer av prosjektgruppen i alfabetisk rekkefølge; (øvrige) medlemmer av ansvarlig faggruppe/hovedkomité/godkjenningsgruppe i alfabetisk rekkefølge; leder/nestleder av ansvarlig faggruppe/hovedkomité/godkjenningsgruppe) (2021). Assessment and updating of a fortification model. Scientific Opinion of the Panel on Nutrition, Dietetic products, Novel food and Allergy of the Norwegian Scientific Committee for Food and Environment. VKM Report 2021:XX, ISBN: 978-82-xxxx-xxx-x, ISSN: 2535-4019. Norwegian Scientific Committee for Food and Environment (VKM), Oslo, Norway.

VKM Report 2021: XX

1 **Assessment and updating of a fortification model**

2 **Preparation of the opinion**

3 The Norwegian Scientific Committee for Food and Environment (Vitenskapskomiteen for mat
4 og miljø, VKM) appointed a project group to draft the opinion. The project group consisted of
5 one VKM member and two VKM staff. The Committee, by the Panel on nutrition, dietetic
6 products, novel food and allergy assessed and approved the final opinion. Before the panel's
7 final approval VKM arranged a public consultation of the opinion from 28. April to 12. May
8 2021.

9 The public consultation was announced on www.vkm.no and in mail to specific recipients.
10 The list of specific recipients was derived from a list of stakeholders from the Norwegian
11 Food Safety Authority), EFSA Focal Point and receivers suggested in the communication plan
12 for VKMs opinion on *Benefit and risk assessment of iodization of household salt and salt used*
13 *in bread and bakery products*. The results of the public consultation on the draft assessment
14 and updating of a fortification model will be published on www.vkm.no in a separate
15 document.

16 **Authors of the opinion**

17 The authors have contributed to the opinion in a way that fulfils the authorship principles of
18 VKM (VKM, 2019). The principles reflect the collaborative nature of the work, and the
19 authors have contributed as members of the project group and/or the VKM Panel on
20 nutrition, dietetic products, novel food and allergy.

21 **Members of the project group** (in alphabetical order after chair of the project group):

22 Lene Frost Andersen – Chair of the project group. Vice-Chair of the Panel on nutrition,
23 dietetic products, novel food and allergy in VKM. Affiliation: 1) VKM; 2) University of Oslo,
24 Department of Nutrition, Institute of Basic Medical sciences.

25 Inger Therese L. Lillegaard – VKM staff. Affiliation: VKM

26 Bente Mangschou – Project leader, VKM staff. Affiliation: VKM.

27 **Members of the Panel on nutrition, dietetic products, novel food and allergy** (in 28 alphabetical order before chair of the Panel):

29 Tor A. Strand - Chair of the Panel on Nutrition, Dietetic Products, Novel Food and Allergy in
30 VKM. Affiliation: 1) VKM; 2) Centre for International Health, University of Bergen and
31 Innlandet Hospital Trust, Department of Research, Lillehammer.

32 Lisbeth Dahl – Member Panel on Nutrition, Dietetic Products, Novel Food and Allergy in VKM.
33 Affiliation: 1) VKM; 2) Institute of Marine Research (IMR), Bergen.

34 Knut Tomas Dalen – Member of the Panel on Nutrition, Dietetic Products, Novel Food and
35 Allergy in VKM. Affiliation: 1) VKM; 2) University of Oslo, Department of Nutrition, Institute
36 of Basic Medical sciences.

37 Kristin Holvik - Member of the Panel on Nutrition, Dietetic Products, Novel Food and Allergy
38 in VKM. Affiliation: 1) VKM; 2) Norwegian Institute of Public Health, Department of Chronic
39 Diseases and Ageing, Oslo.

40 Martinus Løvik – Member of the Panel on Nutrition, Dietetic Products, Novel Food and Allergy
41 in VKM. Affiliation: 1) VKM; 2) Norwegian University of Science and Technology (NTNU),
42 Department of Clinical and Molecular Medicine, Trondheim.

43 Tonje H. Stea - Member of the Panel on Nutrition, Dietetic Products, Novel Food and Allergy
44 in VKM. Affiliation: 1) VKM; 2) University of Agder, Department of Health and Nursing
45 Science.

46 Stine Marie Ulven – Member of the Panel on Nutrition, Dietetic Products, Novel Food and
47 Allergy in VKM. Affiliation: 1) VKM; 2) University of Oslo, Department of Nutrition, Institute
48 of Basic Medical sciences.

49 **Acknowledgement**

50 VKM would like to thank Tue Christensen and co-workers for providing data for vitamin K1
51 and chromium from the Danish National Survey of Diet and Physical Activity (DANSDA) 2011-
52 2013. VKM would also like to thank Inger Ottestad and colleagues for use of the Skedsmo
53 study among home living elderly.

54 **Competence of VKM experts**

55 Persons working for VKM, either as appointed members of the Committee or as external
56 experts, do this by virtue of their scientific expertise, not as representatives for their
57 employers or third-party interests. The Civil Services Act instructions on legal competence
58 apply for all work prepared by VKM.

59

60 Table of Contents

61	Abbreviations	6
62	Background as provided by the Norwegian Food Safety Authority	7
63	Terms of reference as provided by the Norwegian Food Safety Authority	9
64	1 Assessment and revision of the fortification model	11
65	1.1 The fortification model formula	11
66	1.2 Vitamins and minerals included in this revision of the model.....	12
67	1.3 UL/GL values	13
68	1.4 Intake calculations	14
69	1.4.1 Food supplements	16
70	1.4.2 Intake of vitamins and minerals from special products.....	16
71	1.5 Results of the calculations	17
72	1.6 Limitations of the model	23
73	2 Uncertainties	24
74	3 The need for monitoring	25
75	4 Answers to the terms of reference	26
76	5 Data gaps	27
77	References	28
78	Appendix I	32
79	Tolerable Upper Intake Levels (ULs), Safe Upper Levels (SULs) and Guidance Levels (GLs)	
80	used for this fortification model	32
81	Appendix II	35
82	All calculations in the fortification model.....	35
83	Intakes in the 95th percentile in all included age groups	35
84	Intakes in the 90th percentile in all included age groups	41
85	Intakes in the 75th percentile in all included age groups	47
86		

87 Abbreviations

- 88 ALA - acceptable level of addition
- 89 DANSDA - Danish National Survey of Diet and Physical Activity
- 90 EFSA - European Food Safety Authority
- 91 EI95 - the 95th percentile energy intake
- 92 EVM – Expert Group on Vitamins and minerals (UK)
- 93 GL – guidance level
- 94 KBS - kostberegningssystemet
- 95 IOM – Institute of Medicine (US)
- 96 MA – maximal allowance
- 97 NASEM - National Academies of Sciences, Engineering, and Medicine (US)
- 98 NNR – Nordic nutrition recommendations
- 99 OIM – observed individual means
- 100 PFF - proportion of energy from fortified foods
- 101 SUL – safe upper level
- 102 UL – tolerable upper intake level
- 103 VKM – Vitenskapskomiteen for mat og miljø (Norwegian Scientific Committee on Food and
- 104 Environment)

- 105

106 Background as provided by the 107 Norwegian Food Safety Authority

108 According to Norwegian food legislation, foods and drinks fortified with vitamins and
109 minerals could until January 1. 2020 not be placed on the market without special permission
110 from the Norwegian Food Safety Authority (NFSA). This was regulated in the supplementary
111 Norwegian provisions on the addition of vitamins, minerals and certain «other substances» in
112 the regulation (EF) 1925/2006.

113 In 2006 the Norwegian Scientific Committee for Food and Environment adapted a Danish
114 model for assessing applications concerning food fortification with vitamins and minerals into
115 Norwegian conditions. The NFSA also requested the Norwegian Scientific Committee for Food
116 and Environment to assess whether the model can represent a safe assessment of
117 fortification. The Norwegian Scientific Committee for Food and Environment updated the
118 model in 2009¹ and in 2013², and the model was used in the NFSA as a tool in the
119 management of applications on food fortification.

120 From January 1. 2020, amendments regarding the national provisions on the addition of
121 vitamins, minerals and «other substances» entered into force and two so-called positive lists
122 were introduced in the regulation [Forskrift om tilsetning av vitaminer, mineraler og visse
123 andre stoffer til næringsmidler - Lovdata](#), one for the addition of vitamins and minerals, and
124 the other concerning the addition of certain “other substances”.

125 This implies that food business operators may add vitamins and minerals that appear in the
126 positive list to relevant food categories (i.e. in accordance with the requirements in Annex 1
127 in the regulation).

128 If a food business operator wants to add vitamins and minerals to foods in different
129 quantities or to other categories of food from those stated in the positive list, the food
130 business operator must notify the NFSA according to the relevant procedures described in
131 the regulation. As part of the basis to assess these notifications, the NFSA requests an
132 updated model. The model will be fully explained below.

133

¹ [VKM. Modell for vurdering av berikingsaker - revidert versjon 25. februar 2009](#)

² [Revurdering av modell for beriking av mat og drikke - Vitenskapskomiteen for mat og miljø](#)

134 **Thiamine, riboflavin, vitamin B₁₂, biotin, pantothenic acid, potassium,**
135 **chloride and sodium**

136 The addition of thiamine, riboflavin, vitamin B₁₂, biotin, pantothenic acid, potassium, chloride
137 and sodium is excepted from the obligation of notification, provided that the addition not
138 represents a health risk, with reference to footnote 1 in Annex 1 in [Forskrift om tilsetning av](#)
139 [vitaminer, mineraler og visse andre stoffer til næringsmidler - Lovdata](#)

140 **Iodine**

141 Currently there are several food categories in Annex 1 in [Forskrift om tilsetning av vitaminer,](#)
142 [mineraler og visse andre stoffer til næringsmidler - Lovdata](#) where iodine may be added_ i.e.
143 «vegetarian alternatives to milk based drinks», «carbohydrate-electrolyte drinks»,
144 «carbohydrate-electrolyte drinks also containing protein/fat», «meal replacements for weight
145 control», «junior milk», «bars etc.». The NFSA therefore requests that iodine is included in
146 the fortification model as it can be used as a basis to assess future notifications on the
147 addition of iodine.

148

149 Terms of reference as provided by the 150 Norwegian Food Safety Authority

151 The Norwegian Scientific Committee for Food and Environment is requested to update the
152 model for food fortification. The updating of the model should be based upon data from the
153 following food consumption surveys: Ungkost 3 (2015), Småbarnskost 3 (2015) and
154 Spedkost 3 (2020). The updating should include new calculations for energy and vitamins
155 and minerals from the diet and from food supplements.

156 The following is requested included as part of updating the model

- 157 - assessment of the consequences of the amended classification of products such as
158 gluten free foods, sport products, meal replacements for weight control³ etc.
 - 159 ○ to which degree are these products included in the fortification model, and
160 could the intake of vitamins and minerals from these products, in the future,
161 be included in the model?
- 162 - if sufficient scientific data, - assessment of possible inclusion of the additional
163 vitamins and minerals in the model: vitamin K, chromium, fluoride, iodine,
164 molybdenum, manganese, phosphorus (with reference to relevant assessments from
165 the Norwegian Scientific Committee for Food and Environment)
- 166 - inclusion of the recommended Tolerable Upper Intake Levels (UL-levels) from
167 relevant assessments from the Norwegian Scientific Committee for Food and
168 Environment (published after 2013)
- 169 - if available data – inclusion of the age group «70 years and older»
- 170 - a presentation of various versions of the model, based on assumptions that 5%,10
171 %, 15%, 20 % and 25 % of the energy in the diet is derived from fortified foods
- 172 - inclusion of the estimated intake of energy and nutrients from the 95th percentile
173 from foods and drinks in each age group

174 Additionally, if sufficient scientific data:

- 175 - is there a risk that healthy persons (children/adults) on a gluten free diet can exceed
176 UL/GL⁴ for any nutrients?

177 The Norwegian Food Safety Authority requests that the following dietary surveys are used:

- 178 - Spedkost 3 (2020)
- 179 - Småbarnskost 3 (2020)

³ The products were until July 2016 classified as foods for particular nutritional uses and are now classified as regular foods.

⁴ GL: Guidance Level

- 180 - Ungkost 2015
- 181 - Norkost 3 2010-2011
- 182 - and if available: relevant intake data from other Nordic countries or other European
- 183 countries etc.

184 1 Assessment and revision of the 185 fortification model

186 **1.1 The fortification model formula**

187 This is an assessment and revision of previous fortification models by (VKM, 2006; VKM,
188 2013a). It is based on a model suggested by Danish authorities in 2006 (Rasmussen et al.,
189 2006), but some adaptations have been made.

190 The formula for the model is as follows:

$$191 \text{ ALA} = \frac{\text{MA}}{\text{EI}_{95} \times \text{PFF}_n}$$

192

193 *ALA*: acceptable level of addition

194 *MA* (maximal allowance for intake of micronutrients from fortified foods) = (UL, SUL or GL) –
195 (intake from food for the 95th percentile level + mean intake from food supplements among
196 food supplement users).

197 *EI₉₅*: the 95th percentile energy intake

198 *PFF_n*: proportion of energy from fortified foods.

199 UL: Tolerable Upper Intake Level.

200 SUL: Safe Upper Level. Comparable to UL. Term used by the Expert group on Vitamins and
201 Minerals, UK (EVM, 2003).

202 GL: Guidance Level is a value for upper intake in cases where the UL has not been set (see
203 below).

204 The fortification model is based on the intake of vitamins and minerals and energy from the
205 diet at the 95th percentile level in various age groups. As for the additional intakes of
206 vitamins and minerals from food supplements, the mean values among those who have used
207 such supplements are added.

208 As additional information, VKM has calculated intake of vitamins and minerals and energy
209 from the diet at the 90th and 75th percentile levels in various age groups.

210 The total intake from regular foods plus supplements is deducted from the UL/GL for the
211 relevant age group, giving the maximum amount of nutrients that can be “allocated” for food

212 fortification. The maximum amount of a nutrient that can be “allocated” is distributed over
213 the energy intake at the 95th percentile level for the same age group. Based on these
214 inputs, an estimate is made showing which age group is most likely to have an excessive
215 intake of each nutrient.

216 VKM does not have access to any information on how much of present energy intakes that
217 are derived from fortified foods or drinks at an individual level. VKM has calculated for the
218 assumptions that 5, 10, 15, 20 or 25% of the energy intakes are derived from fortified
219 foods/drinks.

220 Finally, the accepted level of addition of vitamins or minerals in the most sensitive age group
221 is listed in Table 1.5-6. Most often, the most sensitive age groups are children between the
222 ages of one and four.

223 This fortification model does not apply to non-caloric products such as e.g. salt or “light
224 products”. “Light products” should be assessed in the same manner as their analogous “non-
225 light products”.

226 **1.2 Vitamins and minerals included in this revision of the model**

227 In this revision of the fortification model, the Norwegian Food Safety Authorities have
228 requested VKM to evaluate maximum limits for addition of the following nutrients: retinol,
229 beta-carotene, vitamin D, vitamin E, niacin, vitamin B6, folic acid, vitamin C, vitamin K,
230 calcium, magnesium, iron, zinc, copper, selenium, iodine, phosphorus, chromium,
231 molybdenum, manganese and fluoride. Vitamin K, iodine, phosphorus, chromium,
232 molybdenum, manganese and fluoride were not included in previous versions of this
233 fortification model from VKM.

234 As thiamine, riboflavin, vitamin B12, biotin, pantothenic acid, potassium, chloride and sodium
235 may be added to several food groups without specified maximum limits, the Norwegian Food
236 Safety Authorities have omitted these substances from the terms of reference. However, as
237 thiamine, riboflavin and vitamin B12 are quite commonly added to foods, VKM has decided to
238 include thiamine, riboflavin and vitamin B12 in the model. VKM have been requested to
239 conduct a benefit and risk assessment of potassium in salt replacers, and will include
240 potassium in the model after finishing the thorough work on this mandate.

241 For vitamin K, chromium, molybdenum and fluoride no intake estimations could be done
242 from national dietary surveys due to lack of concentration data in food and drinks in the
243 Norwegian Food Composition Table. VKM have investigated the possibility of using data from
244 exposure calculations from EFSA or Danish DANSDA (the Danish National Survey of Diet and
245 Physical Activity), and for vitamin K and chromium we have included data from DANSDA
246 2011-2013 for age groups at or above 4 years. It should be noted that data on vitamin K and
247 chromium is not available for 1- and 2-year olds.

248 In 2018, VKM concluded that any dose of manganese as an ingredient in food supplements
249 may be associated with increased risk of negative health effects (VKM, 2018b). This
250 conclusion also applies for manganese as fortification ingredient.

251 **1.3 UL/GL values**

252 Health risks associated with high intakes of vitamins and minerals are described in
253 assessments of Tolerable Upper Intake Levels (ULs) for Foods by the European Food Safety
254 Authority (EFSA), Institute of Medicine/National Academies of Sciences, Engineering, and
255 Medicine (NASEM) (US), Nordic Nutrition Recommendations (NNR) and in previous VKM
256 opinions, and will not be further described in this report.

257 References to the specific ULs/SULs/GLs used for this revised version of the fortification
258 model are given in Appendix I *Tolerable Upper Intake Levels (ULs), Safe Upper Levels (SUL)*
259 *and Guidance Levels (GLs) used for this fortification model*. It should be noted that GLs are
260 based on weaker data than ULs or SULs, and are therefore subject to more uncertainty.

261 The Scientific Committee on Food (SCF) and EFSA has set ULs for the vitamins A, D, E,
262 niacin (nicotinamide and nicotinic acid), vitamin B₆ and folic acid and for the minerals calcium
263 (only for adults), magnesium (only for adults), zinc, copper, iodine and selenium and.
264 Overview table of published ULs from SCF/EFSA is given here.

265 In a series of opinions from VKM on maximum limits for dietary supplements, VKM has also
266 made separate assessments of several ULs based on previous reports not only from
267 SCF/EFSA, but also from IOM/NASEM, NNR or EVM. VKM concluded different from the ULs
268 from SCF/EFSA for vitamin D and children (VKM, 2014), and for zinc (VKM, 2017h). In this
269 fortification model we have used the ULs suggested by VKM for vitamin C, beta-carotene,
270 vitamin D (children and adolescents), calcium (children and adolescents), magnesium
271 (children and adolescents), phosphorus, iron, zinc, chromium and manganese (VKM, 2014,
272 2015a, 2016a, 2016b, 2017b, 2017d, 2017h, 2018a, 2018b).

273 For thiamine we have used temporary GLs suggested by Rasmussen et al., (2006), and for
274 riboflavin and vitamin B12 we have used GLs suggested by Rasmussen et al. (2006), and an
275 Expert Group on Vitamins and Minerals (EVM, 2003).

276 For vitamin K we have used GLs for phylloquinone suggested by (EVM, 2003; VKM, 2018c).

277 Some of the substances do not have a suggested UL for all age groups. The VKM has
278 extrapolated values for the various age groups from values for adults (UL or GL) on the basis
279 of body weight or body surface area (body weight ratio^{0.75}) (Przyrembel, 2018).

280 The UL for vitamin A includes retinol and retinyl esters in µg retinol equivalents per day. The
281 UL for adults for vitamin A does not apply for postmenopausal women.

282 The ULs for beta-carotene, folic acid and magnesium are only applicable for the amount of
283 nutrients added as fortifying agents or in food supplements, and not for those that are
284 naturally present in foods/drinks.

285 **1.4 Intake calculations**

286 The Panel has made use of consumption data from the nationwide Norwegian dietary
287 surveys conducted on various age groups: Adults (Norkost 3, 2010-2011), children and
288 adolescents (Ungkost 3, 2015-2016), 2-year-olds (Småbarnskost, 2019) and infants 12
289 months (Spedkost 3, 2020). Norkost 3 covers only adults up to 70 years. No national dietary
290 survey provides data for persons above 70 years. For adults above 70 years, we have used
291 data from the Skedsmo study (Ottestad et al., 2018). For vitamin K and chromium, we have
292 used data from the Danish National Survey of Diet and Physical Activity (DANSDA, 2011-
293 2013) for 4-, 9- and 13-year olds and adults.

294 1-year-old children: the exposure in 1-year-olds was estimated from Spedkost 3, conducted
295 in 2019 (Paulsen et al., 2020). Spedkost 3 was carried out by the University of Oslo and the
296 Norwegian Institute of Public Health. The food consumption survey was based on a semi-
297 quantitative food frequency questionnaire (FFQ). The caretaker was asked to have the last
298 two weeks in mind when answering the questionnaire. In addition to predefined household
299 units, food amounts were also estimated from photographs. A total of 1957 1-year-olds
300 participated. The participation rate was 65%. Only children that did not receive breastmilk
301 were included in this assessment (n=1024).

302 2-year-old children: The intake in 2-year-olds was estimated from Småbarnskost 3,
303 conducted in 2019 (Astrup et al., 2020). Småbarnskost 3 was carried out by the University of
304 Oslo and the Norwegian Institute of Public Health. This food consumption survey was based
305 on a semi-quantitative food frequency questionnaire (FFQ). The caretaker was asked to have
306 the last two weeks in mind when answering the questionnaire. In addition to predefined
307 household units, food amounts were also estimated from photographs. A total of 1413 2-
308 year-olds participated. The participation rate was 47%.

309 4-, 9- and 13-year-old children/adolescents: The intake in 4-, 9- and 13-year-old children
310 was estimated from the national food consumption survey Ungkost 3 (Hansen et al., 2016;
311 Hansen et al., 2015). The Ungkost 3 study was carried out by the University of Oslo, the
312 Norwegian Food Safety Authority, the Norwegian Directorate of Health and the Norwegian
313 Institute of Public Health in 2015 for 4th and 8th graders (8-9-year-olds and 12-13-year-
314 olds), and in 2016 for 4-year-old children. The dietary assessment tool was a 4 days
315 validated web-based food diary. A total of 399 4-year-old, 636 9-year-old and 687 13-year-
316 old children participated. The participation rates were 20, 55 and 53%, respectively.

317 Adults 18-70 years: Norkost 3 was based on two 24-hour recalls by telephone at least one
318 month apart. Food amounts were presented in household measures or estimated from
319 photographs (Totland et al., 2012). The Norkost 3 study was carried out by the University of

320 Oslo, the Norwegian Food Safety Authority, the Norwegian Directorate of Health and the
321 Norwegian Institute of Public Health. A total of 1787 adults aged 18-70 years participated.
322 The participation rate was 37%.

323 Adults 70 years and above: A dietary study in adults living in the municipality of Skedsmo
324 were used for the age 70 and above (Ottestad et al., 2018). The dietary assessment tool
325 was two 24-hour recalls, the first face-to-face, and the second was an unannounced
326 telephone interview 14 days, or more, later. Portion sizes were estimated in household
327 measures, and with the same type of photographs as in Norkost 3. A total of 387 adults
328 were included in the cross-sectional dietary survey. The participation rate was 15%.

329 DANSDA 2011-2013: Intake of food and drink was collected from a representative sample of
330 3946 individuals aged 4-75 years with a participation rate of 54% (Pedersen et al., 2015).
331 The participants kept a food record for seven consecutive days in a pre-coded (semi-closed)
332 questionnaire with answering categories for the most commonly consumed foods and dishes
333 in the Danish diet. The participants in the results presented in this report were grouped (4 to
334 9-years-olds, 10 to 17-years-olds and 18 to 75-years-olds).

335 The intake of nutrients from fortified foods and drinks that are already available on the
336 Norwegian market are included only to a limited extent in the calculations. The intake of
337 vitamins A and D from fortified butter, margarine and milk for which general permission has
338 been granted by the Norwegian Food Safety Authority are included in our exposure
339 estimates, whereas no other fortifications are included.

340 The exposure estimates are based on means of survey days (OIMs) from Norkost 3, Ungkost
341 3, DANSDA and the Skedsmo study, whereas data for 1- and 2-year-olds are based on FFQ.

342 VKM has estimated the intake of each included nutrient/substance in the 95th percentile
343 from regular foods and drinks (not including food supplements) in each age group. The 95th
344 percentile is commonly used in risk assessments, and covers most of the consumers. It
345 should be noted that there is a higher uncertainty associated with the 95th percentile than
346 the mean value, and the 95th percentile is more likely an overestimation than
347 underestimation of intake.

348 With use of the dietary and supplement intake calculations the amount of nutrients that can
349 be allocated for fortification have been estimated. Tables 1.5-6 accounts for the most
350 vulnerable groups, i.e. the age group that tolerates the lowest addition per 100 kcal before a
351 risk of exceeding the UL/SUL/GL arises. The age groups that (at the 95th percentile level)
352 already have an intake from diet and food supplements that exceeds the UL/GL will have a
353 tolerance for fortification after the calculations that is below zero (a negative number). In
354 practice this means that fortification with this nutrient cannot be done without exceeding
355 UL/SUL/GL. The negative numbers have been changed to zero in the column on the extreme
356 right in Table 1.5-6.

357 **1.4.1 Food supplements**

358 The use of food supplements in Norway is common among adults (47% among the men,
359 and 58% among the women) (Totland et al., 2012). In children and adolescents, the intake
360 of food supplements range from 43% among the 13-year olds to 79% among breastfed 1-
361 year-olds (67% in non-breastfed).

362 In the fortification model, the mean values of nutrient intake from food supplements (users
363 only) is added to the intakes from regular foods and drinks to estimate the total intake of
364 nutrients.

365 As most of the calculations are based on mean intake among food supplement users, the
366 estimates do not account for high intakes from food supplements. There will be uncertainty
367 associated with the estimates for food supplements with few users.

368 **1.4.2 Intake of vitamins and minerals from special products**

369 VKM is requested to comment on to which degree products such as gluten free products,
370 sports products, meal replacers for weight control etc. are included in this fortification model.
371 Additionally, if sufficient scientific data, is there a risk that healthy persons (children/adults)
372 on a gluten free diet can exceed UL/GL for any nutrients?

373 These products are only to a very limited extent included in the fortification model. These
374 products are registered in the food surveys for adults (2x24 hours recall) and children and
375 adolescents (4 days food diary) if they have been consumed by some of the participants.
376 However, very few people in the surveys have eaten such products.

377 In the dietary surveys for 1- and 2-year-olds (FFQs) fortified infant formula and baby
378 porridges are included, whereas gluten free products are not. However, both the gluten
379 containing and the gluten-free alternatives are subject to the same regulation for addition of
380 vitamins and minerals.

381 VKM received data on content of vitamins and minerals in several gluten free products, such
382 as, bread, crisp breads and breakfast cereals, see Table 1.4.2-1. Most products contain
383 vitamins and minerals in similar concentrations as their gluten containing alternatives. Only
384 two of the products VKM received data on, the gluten free Cornflakes from AS Nestle Norge,
385 contained vitamins and minerals in high concentrations, especially of iron, niacin and folic
386 acid. Individuals consuming these products will be at risk of exceeding the ULs/GLs for iron,
387 niacin and folic acid.

388 **Table 1.4.2-1** Concentration of some nutrients per 100 g in gluten free products available in Norway.
389 Product names in Norwegian. Data provided by the Norwegian Food Safety Authority.

	Niacin	Vitamin B6	Folic acid	Thiamine	Riboflavin	Vitamin B12	Iron	Magnesium
AS Nestle Norge								

	Niacin	Vitamin B6	Folic acid	Thiamine	Riboflavin	Vitamin B12	Iron	Magnesium
<i>Breakfast cereals</i>								
GoFree Cornflakes								121 mg
GoFree Chocolate Cornflakes	16 mg	1.4 mg	183 µg					106 mg
Semper								
<i>Cake-/flour mix</i>								
Mix Glutenfri	4 mg	0.35 mg	40 µg	0.4 mg	0.4 mg		6.5 mg	
Glutenfri Mix med fiber	4 mg	0.35 mg	40 µg	0.4 mg	0.4 mg		8.0 mg	
Fin Mix	4 mg	0.35 mg	40 µg	0.4 mg	0.4 mg		7.5 mg	
Grov Mix	4 mg	0.35 mg	40 µg	0.4 mg	0.4 mg		8.0 mg	
<i>Crispbread</i>								
Rosmarinknekkebrød	4 mg	0.3 mg	40 µg	0.3 mg	0.3 mg		12 mg	
Surdeigsknekkebrød	4 mg	0.3 mg	40 µg	0.3 mg	0.3 mg		8.5 mg	
<i>Breads</i>								
Trio rundstykker	2.4 mg	0.21 mg	30 µg	0.20 mg	0.21 mg		4.5 mg	
Toasty fiber	1.8 mg	0.14 mg	27 µg	0.17 mg	0.21 mg		4.0 mg	
Mykt fjellbrød	2.4 mg	0.21 mg	30 µg	0.2 mg	0.21 mg		5.3 mg	
Minibaguetter	2.7 mg	0.21 mg	30 µg	0.2 mg	0.21 mg		4.5 mg	
Grova grovbrød	1.8 mg	0.14 mg	27 µg	0.12 mg	0.12 mg		4.0 mg	
Minibaguetter fiber	2.7 mg	0.21 mg	30 µg	0.2 mg	0.21 mg		4.5 mg	
<i>Breakfast cereals</i>								
Special flakes	25 mg	2.5 mg	250 µg	1.8 mg	2.0 mg	1.3 µg	20 mg	
Flakes & Red berries	25 mg	2.5 mg	250 µg	1.8 mg	2.0 mg	1.3 µg	20 mg	
<i>Pizza</i>								
Pizzabunn	2.4 mg	0.21 mg	30 µg	0.2 mg	0.21 mg		4.5 mg	

390

391 1.5 Results of the calculations

392 All the calculations are given in Appendix II *All calculations in the fortification model*.

393 All calculations have been made according to the model formula – with Norwegian dietary
394 data and Norwegian data for the intake of food supplements, except for vitamin K1 and
395 chromium in which Danish data was used. In addition to the requested calculations for
396 intakes in the 95th percentile, we have made similar estimates for the 90th and 75th
397 percentiles.

398 A summary of the calculations can be viewed in the Summary Tables 1.5-1 to 1.5-5 below.
399 The column to the extreme right in these tables shows the most vulnerable and limiting age
400 group. Table 1.5-6 shows the amount of the various nutrients that can be added to
401 foodstuffs in Norway (per 100 kcal) without exceeding UL/GL in 95th percentile of the
402 population at the various age groups if 25, 20, 15, 10 or 5% of the energy in the diet is
403 derived from fortified products if all the requested age groups are included. The negative
404 numbers in Tables 1.5-1 to 1.5-5 have been changed to zero in Table 1.5-6. As can be seen

405 from the results in these Summary Tables, the assumptions of 5 or 25% of the energy
 406 derived from fortified foods affect the amount that may be added per 100 kcal, but does not
 407 affect which nutrients that could be added to foods or drinks. Inclusion of the youngest age
 408 groups affects both which nutrients that may be added (especially for vitamin D and several
 409 of the minerals or trace elements) and the amount that could be added without exceeding
 410 the ULs/GLs.

411 Example if 25 or 5% of the energy is derived from fortified foods/drinks:

412 The most vulnerable group for vitamin D intake is 1-year-old children. For this group, the
 413 95th percentile intake of vitamin D from food is 16.8 µg per day, and the intake from food
 414 supplements is 9.6 µg, giving a total possible intake of 26.4 µg vitamin D/day. The UL for
 415 this age group is 25 µg, i.e. an MA of -1.4 µg. EI95 is 2000 kcal/day for this age group.

416 For 1-year-olds $ALA_{25} = [-1.4 \mu\text{g} / (2000 \text{ kcal} \times 0.25)] \times 100 = -0.3 \mu\text{g}$ vitamin D per 100 kcal
 417 (changed to zero in the column on the extreme right in Table 1.5-6)

418 For 1-year-olds $ALA_5 = [-1.4 \mu\text{g} / (2000 \text{ kcal} \times 0.05)] \times 100 = -1.4 \mu\text{g}$ vitamin D per 100 kcal
 419 (changed to zero in the column on the extreme right in Table 1.5-6)

420 For men 18-70 years $ALA_{25} = [69.2 \mu\text{g} / (4200 \text{ kcal} \times 0.25)] \times 100 = 7 \mu\text{g}$ vitamin D per 100
 421 kcal

422 **Table 1.5-1** Summary Table 25 E%. Amount of nutrients that can be added per 100 kcal of foodstuff
 423 for each included age group according to the calculations in the fortification model assuming that 25%
 424 of the energy in the diet is derived from fortified foods.

E%: 25	Men, 70+	Women, 70+	Men 18-70 y	Women, 18-70 y	Adolescents, 13 y	Children, 9 y	Children, 4 y	Children, 2 y	Infants, 1 y	Most sensitive age group
Retinol, µg	101	207	80	183	42	25	-34	-268	-362	1-year-olds
β-carotene, µg	531	616	244	337	387	320	350	200	200	1- and 2-year-olds
Vitamin D, µg	9	11	7	11	4	1.5	2	1.4	-0.3	1-year-olds
Tocopherol, mg	37	44	25	37	25	22	20	16	14	1-year-olds
Thiamine, mg	6	8	4	6	4	4	4	3	2	1-year-olds
Riboflavin, mg	5	6	4	5	3	3	3	2	2	1- and 2-year-olds
Niacin, mg	118	138	80	118	60	51	38	25	24	1-year-olds
Vitamin B12, µg	266	317	188	272	170	158	160	104	105	2-year-olds
Vitamin B6, mg	3	3	1.9	2.8	2.2	1.2	0.9	0.5	0.5	1- and 2-year-olds
Folic acid, µg	116	144	79	117	64	53	49	25	22	1-year-olds
Vitamin C, mg	103	117	58	88	133	167	97	45	29	1-year-olds
Calcium, mg	126	164	28	100	69	178	179	135	178	Adult men
Magnesium, mg	11	32	16	19	23	31	23	5	4	1-year-olds
Iron, mg	4	5	1.8	2.2	1.1	0.3	-1.6	-1.8	-5.2	1-year-olds
Zinc, mg	2	3	0.7	2.1	0.3	0.5	-1.0	-2.1	-2.5	1-year-olds

E%: 25	Men, 70+	Women, 70+	Men 18-70 y	Women, 18-70 y	Adolescents, 13 y	Children, 9 y	Children, 4 y	Children, 2 y	Infants, 1 y	Most sensitive age group
Copper, mg	0.3	0.4	0.1	0.3	0.3	0.2	0	-0.1	-0.1	1- and 2-year-olds
Selenium, µg	19	28	12	23	16	9	2	-3.4	-3.6	1-year-olds
Iodine, µg	-14	19	-8	12	20	2,2	-2	-30	-34	1-year-olds
Phosphorus, mg	85	159	-27	88	-291	-319	-433	-387	-330	4-year-olds
Vitamin K1, µg	NA	NA	83	112	69	62	42	NA	NA	4-year-olds
Chromium, µg	NA	NA	2145	2887	1620	1059	1059	NA	NA	4- and 9-year-olds

425 NA: not available

426 **Table 1.5-2** Summary Table 20 E%. Amount of nutrients that can be added per 100 kcal of foodstuff
427 for each included age group according to the calculations in the fortification model assuming that 20%
428 of the energy in the diet is derived from fortified foods.

E%: 20	Men, 70+	Women, 70+	Men 18-70 y	Women, 18-70 y	Adolescents, 13 y	Children, 9 y	Children, 4 y	Children, 2 y	Infants, 1 y	Most sensitive age group
Retinol, µg	126	259	100	229	53	32	-43	-335	-452	1-year-olds
β-carotene, µg	664	770	304	421	484	400	438	250	250	1 and 2 year olds
Vitamin D, µg	11	14	8	13	5	1.9	2.0	1.7	-0.4	1-year-olds
Tocopherol, mg	46	54	31	47	32	28	25	20	18	1-year-olds
Thiamine, mg	8	9	5	6	5	5	5	3	3	1- and 2-year-olds
Riboflavin, mg	7	8	5	5	4	4	4	2	2	1- and 2-year-olds
Niacin, mg	148	172	100	148	75	64	48	32	30	1-year-olds
Vitamin B12, µg	333	397	235	341	213	198	200	130	131	2-year-olds
Vitamin B6, mg	4	4	2.4	3.5	2.7	1.5	1.1	0.7	0.6	1-year-olds
Folic acid, µg	145	180	99	147	80	66	61	31	28	1-year-olds
Vitamin C, mg	128	146	72	110	166	209	121	56	36	1-year-olds
Calcium, mg	158	205	36	125	86	222	224	169	223	Adult men
Magnesium, mg	14	40	20	24	29	38	29	6.3	4.9	1-year-olds
Iron, mg	4	6	2.3	2.8	1.4	0.3	-2.1	-2.3	-6.5	1-year-olds
Zinc, mg	2	4	0.9	2.6	0.3	0.6	-1.3	-2.6	-3.1	1-year-olds
Copper, mg	0,4	0,5	0.2	0.3	0.4	0.3	-0.1	-0.1	-0.2	1-year-olds
Selenium, µg	24	35	15	29	20	11	3,0	-4.3	-4.5	1-year-olds
Iodine, µg	-17	24	-10	14	25	3	-3	-37	-43	1-year-olds
Phosphorus, mg	107	199	-33	110	-364	-399	-541	-484	-412	4-year-olds
Vitamin K1, µg	NA	NA	103	140	86	78	53	NA	NA	4-year-olds
Chromium, µg	NA	NA	2681	3609	2025	1324	1324	NA	NA	4- and 9-year-olds

429 NA: not available

430 **Table 1.5-3** Summary Table 15 E%. Amount of nutrients that can be added per 100 kcal of foodstuff
 431 for each included age group according to the calculations in the fortification model assuming that 15%
 432 of the energy in the diet is derived from fortified foods.

E%: 15	Men, 70+	Women, 70+	Men 18-70 y	Women, 18-70 y	Adolescents, 13 y	Children, 9 y	Children, 4 y	Children, 2 y	Infants, 1 y	Most sensitive age group
Retinol, µg	168	345	134	306	70	42	-57	-446	-603	1-year-olds
β-carotene, µg	885	1027	406	562	645	533	583	333	333	1 and 2 year olds
Vitamin D, µg	15	19	11	18	6,7	2.5	2.6	2.3	-0.5	1-year-olds
Tocopherol, mg	61	73	42	62	42	37	33	26	24	2-year-olds
Thiamine, mg	11	13	7	11	7	6	7	4	4	1- and 2-year-olds
Riboflavin, mg	9	11	6	9	6	5	5	3	3	1- and 2-year-olds
Niacin, mg	197	230	133	197	99	85	64	42	41	1-year-olds
Vitamin B12, µg	444	529	313	454	284	264	267	174	174	1- and 2-year-olds
Vitamin B6, mg	5	6	3.2	4.7	3.7	2.0	1.4	0.9	0.8	1-year-olds
Folic acid, µg	193	241	131	196	107	89	82	41	37	1-year-olds
Vitamin C, mg	171	195	96	146	222	279	162	74	48	1-year-olds
Calcium, mg	211	273	47	166	115	296	299	225	297	Adult men
Magnesium, mg	18	53	27	32	38	51	38	8.3	6.5	1-year-olds
Iron, mg	6	9	3.0	3.7	1.8	0.5	-2.7	-3.0	-8.7	1-year-olds
Zinc, mg	3	5	1.2	3.4	0.5	0.8	-1.7	-3.5	-4.1	1-year-olds
Copper, mg	0,5	0,7	0.2	0.5	0.5	0.3	-0.1	-0.2	-0.2	1- and 2-year-olds
Selenium, µg	32	47	20	39	26	15	4	-5.7	-6.0	1-year-olds
Iodine, µg	-23	32	-14	19	33	3.7	-4	-50	-57	1-year-olds
Phosphorus, mg	142	265	-44	147	-485	-532	-721	-645	-549	4-year-olds
Vitamin K1, µg	NA	NA	138	187	115	104	71	NA	NA	4-year-olds
Chromium, µg	NA	NA	3575	4812	2699	1765	1765	NA	NA	4- and 9-year-olds

433 NA: not available

434 **Table 1.5-4** Summary Table 10 E%. Amount of nutrients that can be added per 100 kcal of foodstuff
 435 for each included age group according to the calculations in the fortification model assuming that 10%
 436 of the energy in the diet is derived from fortified foods.

E%: 10	Men, 70+	Women, 70+	Men 18-70 y	Women, 18-70 y	Adolescents, 13 y	Children, 9 y	Children, 4 y	Children, 2 y	Infants, 1 y	Most sensitive age group
Retinol, µg	252	517	200	458	105	63	-86	-669	-905	1-year-olds
β-carotene, µg	1328	1540	609	843	968	800	875	500	500	1 and 2 year olds
Vitamin D, µg	22	29	16	26	10	3.8	4.0	3.5	-0.7	1-year-olds
Tocopherol, mg	92	109	62	93	63	56	50	39	36	1-year-olds
Thiamine, mg	16	19	11	16	10	9	10	6	6	1- and 2-year-olds
Riboflavin, mg	13	16	9	13	8	8	8	4	4	1- and 2-year-olds
Niacin, mg	295	345	200	295	149	128	96	64	61	1-year-olds

E%: 10	Men, 70+	Women, 70+	Men 18-70 y	Women, 18-70 y	Adolescents, 13 y	Children, 9 y	Children, 4 y	Children, 2 y	Infants, 1 y	Most sensitive age group
Vitamin B12, µg	666	793	470	681	425	396	401	261	261	1- and 2-year-olds
Vitamin B6, mg	7	9	4.7	7.0	5.5	3.0	2.2	1.3	1.2	1-year-olds
Folic acid, µg	289	361	197	293	161	133	123	62	56	1-year-olds
Vitamin C, mg	257	292	145	220	333	418	243	112	72	1-year-olds
Calcium, mg	316	409	71	250	172	444	448	338	446	Adult men
Magnesium, mg	28	80	41	48	58	76	57	13	10	1-year-olds
Iron, mg	9	13	4.6	5.5	2.8	0.7	-4.1	-4.5	-13.0	1-year-olds
Zinc, mg	5	8	1.8	5.2	0.7	1.2	-2.6	-5.3	-6.2	1-year-olds
Copper, mg	0,7	1,0	0.3	0.7	0.7	0.5	-0.1	-0.3	-0.4	1-year-olds
Selenium, µg	48	71	30	59	39	22	6.0	-8.5	-9.0	1-year-olds
Iodine, µg	-34	47	-20	29	50	5.6	-6.0	-75	-86	1-year-olds
Phosphorus, mg	213	398	-66	221	-728	-798	-	-968	-824	4-year-olds
Vitamin K1, µg	NA	NA	206	280	173	156	106	NA	NA	4-year-olds
Chromium, µg	NA	NA	5363	7218	4049	2648	2648	NA	NA	4- and 9-year-olds

437 NA: not available

438 **Table 1.5-5** Summary Table 5 E%. Amount of nutrients that can be added per 100 kcal of foodstuff
439 for each included age group according to the calculations in the fortification model assuming that 5%
440 of the energy in the diet is derived from fortified foods.

E%: 5	Men 70+	Women, 70+	Men 18-70 y	Women, 18-70 y	Adolescents, 13 y	Children, 9 y	Children, 4 y	Children, 2 y	Infants, 1 y	Most sensitive age group
Retinol, µg	503	1034	401	917	211	126	-172	-1338	-1809	1-year-olds
β-carotene, µg	2655	3080	1218	1686	1935	1600	1750	1000	1000	1 and 2 year olds
Vitamin D, µg	44	57	33	53	20	7.6	7.9	6.9	-1.4	1-year-olds
Tocopherol, mg	184	218	125	186	126	112	99	79	72	1-year-olds
Thiamine, mg	32	38	22	32	20	18	18	13	12	1-year-olds
Riboflavin, mg	27	32	18	27	17	15	13	9	9	1- and 2-year-olds
Niacin, mg	590	689	400	590	298	256	192	127	122	1-year-olds
Vitamin B12, µg	1332	1586	940	1362	851	792	720	521	523	2-year-olds
Vitamin B6, mg	15	17	9.5	14.1	11	6.1	4.3	2.6	2.3	1-year-olds
Folic acid, µg	579	722	394	587	321	266	245	123	111	1-year-olds
Vitamin C, mg	514	584	289	439	666	836	485	223	143	1-year-olds
Calcium, mg	632	819	142	499	344	889	896	675	892	Adult men
Magnesium, mg	55	160	81	96	115	153	114	25	20	1-year-olds
Iron, mg	18	26	9.1	11.0	5.5	1.4	-8.2	-9.0	-26.0	1-year-olds
Zinc, mg	10	16	3.6	10.3	1.4	2.4	-5.1	-11	-12	1-year-olds
Copper, mg	1	2	0.6	1.4	1.4	1.0	-0.2	-0.5	-0.7	1-year-olds
Selenium, µg	95	142	61	117	78	44	12	-17	-18	1-year-olds
Iodine, µg	-69	95	-41	58	100	11	-12	-149	-171	1-year-olds

E%: 5	Men 70+	Women, 70+	Men 18-70 y	Women, 18-70 y	Adolescents, 13 y	Children, 9 y	Children, 4 y	Children, 2 y	Infants, 1 y	Most sensitive age group
Phosphorus, mg	427	796	-133	441	-1456	-1597	-2164	-1935	-1648	4-year-olds
Vitamin K1, µg	NA	NA	413	561	346	312	212	NA	NA	4-year-olds
Chromium, µg	NA	NA	10726	14437	8098	5296	5296	NA	NA	4- and 9-year-olds

441 NA: not available

442 The maximum addition of each nutrient that can be added to 100 kcal of foods or drinks
443 according to this fortification model is given in Table 1.5-6 under the assumptions that 25,
444 20, 15, 10 or 5% of the energy in the diet is derived from fortified foods or drinks and
445 inclusion of all the requested age groups (from 1-year-olds to above 70 years). It should be
446 noted that for vitamin K1 and chromium, data for 1- and 2-year-olds are not available.

447 **Table 1.5-6** Maximum amount of nutrients that can be added per 100 kcal of foodstuff according to
448 the calculations in the fortification model assuming 25, 20, 15, 10 or 5 E% from fortified foods and
449 inclusion of all age groups requested (from 1-year-olds to above 70 years).

Nutrients	25 E%	20 E%	15 E%	10 E%	5 E%
Retinol, µg	0	0	0	0	0
β-carotene, µg	200	250	333	500	1000
Vitamin D, µg	0	0	0	0	0
Tocopherol, mg	14	18	24	36	72
Thiamine, mg	2	3	4	6	12
Riboflavin, mg	2	2	3	4	9
Niacin, mg	24	30	41	61	122
Vitamin B12, µg	104	130	174	261	521
Vitamin B6, mg	0.5	0.6	0.8	1.2	2.3
Folic acid, µg	22	28	37	56	111
Vitamin C, mg	29	36	48	72	143
Calcium, mg	28	36	47	71	142
Magnesium, mg	4	4.9	6.5	10	20
Iron, mg	0	0	0	0	0
Zinc, mg	0	0	0	0	0
Copper, mg	0	0	0	0	0
Selenium, µg	0	0	0	0	0
Iodine, µg	0	0	0	0	0
Phosphorus, mg	0	0	0	0	0
Vitamin K1*, µg	42	53	71	106	212
Chromium*, µg	1059	1324	1765	2648	5296
Manganese, mg	0	0	0	0	0

450 *Data for 1- and 2-year-olds are not available.

451 **1.6 Limitations of the model**

452 The model does not include fortification of foods such as salt, seasonings, water or
453 foodstuffs that do not naturally contain energy.

454 Since light products are to be assessed in the same manner as their analogous non-light
455 products, this leads to a higher content of added nutrients per unit of energy for light
456 products than that which the model has included in the calculations. If light products
457 (including products that provide no energy) become a major source of a nutrient, the
458 prerequisites for the calculations in the model will no longer apply.

459 The model includes mean intakes of vitamins and minerals from supplements (users only).
460 Individuals with higher intakes than the means, may not be sufficiently protected from
461 exceedance of ULs/GLs.

462 Consumers with high energy intakes from a few single food or drink item may not be
463 sufficiently protected from high intakes if their energy sources are fortified above the energy
464 percent given in the model.

465 Nutritional policy authorities require special measures to be implemented for some nutrients
466 in order to increase the intake in certain sections of the population. The model does not
467 assess the risk of a low intake of nutrients (only high intakes) and does not take into account
468 the fact that food fortification targeted directly at certain groups of the population may be
469 desirable e.g. population groups with intakes of iron and vitamin D below recommended
470 levels.

471

472 2 Uncertainties

473 Section 1.6 describes the main limitations in the model itself, e.g. what the model does not
474 include or cover, and certain population group that might not be protected from exceeding
475 ULs/GLs with this model. In this chapter we describe uncertainties underlying the data inputs
476 in the model.

477 Several of the ULs are established based on older studies, and newer published studies
478 should be included. The data for establishing ULs are often scarce, especially in children and
479 adolescents. For nutrients in the model that have been estimated based on GLs the
480 uncertainty is high. Some of the GLs are based on one or few studies and in small population
481 groups not necessarily representative for the whole population.

482 In the model, we have used intakes in the 95th percentiles as estimates for nutrients and
483 energy from foods and drinks. Data in the 95th percentile is generally associated with
484 uncertainty, and the estimated intakes in the 95th percentiles may be higher than the true
485 intakes.

486 For intakes of nutrients from food supplements we have used mean intakes among
487 supplement users of the specific nutrient only. The input data for supplements with rarely
488 included nutrients are associated with uncertainty. For some nutrients, especially in the
489 younger age groups, these means are based on very few individuals e.g. from one to ten
490 subjects.

491 The intake of nutrients from fortified foods and drinks that are already available on the
492 Norwegian market are included only to a limited extent in the calculations. The intake of
493 vitamins A and D from fortified butter, margarine and milk for which general permission has
494 been granted by the Norwegian Food Safety Authority are included in our exposure
495 estimates as are infant formulas and baby porridges, whereas no other fortifications are
496 included. For folic acid, magnesium and beta-carotene, intake data from foods or drinks are
497 not included because the UL for these nutrients are established for supplemental forms only.
498 As we cannot separate natural containing folate, magnesium or beta-carotene from added
499 the added forms in e.g. infant formulas or baby porridges, any contribution to folic acid,
500 magnesium or beta-carotene from infant formulas or baby porridges are not accounted for.

501 The food consumption data used in this assessment are from five different studies with four
502 different dietary assessment methods.

503

504 3 The need for monitoring

505 To ensure that food fortification does not represent a health risk, the prerequisites on which
506 the model is based must be monitored:

- 507 - Surveillance of intake of fortified products on the Norwegian market, including intake
508 of fortified light products and products without energy, e.g. fortified salt, waters etc.
509 Without good knowledge on available fortified products on the market, and good
510 consumption data of these products, all intake estimates for vitamins and minerals
511 will be associated with uncertainty in the whole population.
- 512 - The intake of nutrients and energy from food/drinks and vitamin and mineral
513 supplements in different groups of the population.

514

515 4 Answers to the terms of reference

516 Products such as gluten free products, sports products, meal replacers for weight control etc.
517 are only to a very limited extent included in the fortification model. These products are
518 registered in the food surveys if they have been consumed by some of the participants.
519 However, very few people in the surveys have eaten such products.

520 Due to lack of data on tolerable upper intake levels/ upper guidance levels or exposure data,
521 molybdenum, and fluoride could not be included in the model. For vitamin K1 and chromium
522 exposure data was not available for 1- and 2-year-olds.

523 The tolerable upper intake levels/ safe upper levels/ guidance levels VKM have used for this
524 opinion are given in Appendix I.

525 Maximum amount of nutrients that can be added per 100 kcal of foodstuff according to the
526 calculations in the fortification model assuming 25, 20, 15, 10 or 5 E% from fortified foods is
527 given in Summary Tables 1.5-1 to 1.5-5 and Table 1.5-6. All the calculations are given in
528 Appendix II.

529 In gluten free products that are fortified with nutrients, the concentrations are in
530 approximately the same levels as the gluten containing alternatives

531 Persons (children/adults) on a gluten free diet have therefore generally the same risk of
532 exceeding ULs/GLs as persons that consume the gluten containing products. There are two
533 specific gluten free breakfast cereals on the Norwegian market that contain much higher
534 concentrations of iron, folic acid and niacin (up to ten fold). Subjects consuming high
535 amounts of these specific breakfast cereals will be at risk of exceeding UL for these
536 nutrients.

537 Dietary patterns develop and change over time. To keep up with the changes, the
538 fortification model needs to be updated along with new national dietary surveys.

539

540 5 Data gaps

- 541 • Intake of nutrients derived from fortified products.
- 542 • More knowledge about supplement use, especially in certain age groups such as
- 543 children, adolescents and elderly.
- 544 • Updated ULs for several nutrients with emphasis on studies in the younger age
- 545 groups. Several nutrients lack ULs, and studies investigating negative health effects
- 546 from high intakes are scarce, especially in children and adolescents.
- 547 • Food composition data for vitamin K, chromium, molybdenum, manganese and
- 548 fluoride.
- 549 • Intake of products such as gluten free products, sports products, meal replacers for
- 550 weight control etc.
- 551

552 References

- 553 Astrup H., Myhre J.B., Andersen L.F., Kristiansen A.L. (2020) Småbarnskost 3,
554 Landsomfattende undersøkelse av kostholdet blant 2-åringer i Norge, National
555 Institute of Public Health, Oslo Norway. ISBN: 978-82-8406-100-9.
- 556 EFSA. (2018) Overview on Tolerable Upper Intake Levels as derived by the Scientific
557 Committee on Food (SCF) and the EFSA Panel on Dietetic Products, Nutrition and
558 Allergies (NDA), Summary of Tolerable Upper Intake Levels – version 4 (September
559 2018), European Food Safety Authority.
- 560 EVM. (2003) Safe Upper Levels for Vitamins and Minerals, in: Expert Group on Vitamins and
561 Minerals (Ed.), Food Standard Agency, London, UK.
- 562 Hansen L.B., Myhre J.B., Andersen L.F. (2016) Ungkost 3, Landsomfattende
563 kostholdsundersøkelse blant 4-åringer i Norge, National Institute of Public Health,
564 Oslo, Norway.
- 565 Hansen L.B., Myhre J.B., Johansen A.M.W., Paulsen M.M., Andersen L.F. (2015) Ungkost 3,
566 Landsomfattende kostholdsundersøkelse blant elever i 4. -og 8. klasse i Norge,
567 National Institute of Public Health, Oslo, Norway.
- 568 Ottestad I., Ulven S.M., Oyri L.K.L., Sandvei K.S., Gjevestad G.O., Bye A., Sheikh N.A., Biong
569 A.S., Andersen L.F., Holven K.B. (2018) Reduced plasma concentration of branched-
570 chain amino acids in sarcopenic older subjects: a cross-sectional study. *Br J Nutr*
571 120:445-453. DOI: 10.1017/S0007114518001307.
- 572 Paulsen M.M., Myhre J.B., Andersen L.F., Kristiansen A.L. (2020) Spedkost 3,
573 Landsomfattende undersøkelse av kostholdet blant spedbarn i Norge, 12 måneder,
574 National Institute of Public Health, Oslo, Norway. ISBN: 978-82-8406-099-6.
- 575 Pedersen A.N., Christensen T., Matthiessen J., Knudsen V.K., Rosenlund-Sørensen M., Biloft-
576 Jensen A., Hinsch H.-J., Ygil K.H., Kørup K., Saxholt E., Trolle E., Søndergaard A.B.,
577 Fagt S. (2015) Dietary habits in Denmark 2011-2013, Main results, DTU
578 Fødevareinstituttet, Søborg, Denmark. ISBN: 978-87-93109-39-1.
- 579 Przyrembel H. (2018) 4. Filling the data gaps – extrapolations. 4.1 Extrapolations – allometric
580 or isometric scaling?, in: I. Tetens, et al. (Eds.), *The dual risk approach in nutrition:
581 Present and future perspectives and challenges*, The Nordic Council of Ministers,
582 Copenhagen, Copenhagen, Denmark. pp. 33-34.
- 583 Rasmussen S.E., Andersen N.L., Dragsted L.O., Larsen J.C. (2006) A safe strategy for
584 addition of vitamins and minerals to foods. *European Journal of Nutrition* 45:123-135.
585 DOI: DOI 10.1007/s00394-005-0580-9.
- 586 Totland T.H., Melnæs B.K., Lundberg-Hallén N., Helland-Kigen K.M., Lund-Blix N.A., Myhre
587 J.B., Johansen A.M.W., Løken E.B., Andersen L.F. (2012) Norkost 3, En
588 landsomfattende kostholdsundersøkelse blant menn og kvinner i Norge i alderen 18-
589 70 år, 2010-11, Helsedirektoratet, Oslo, Norway.

- 590 VKM. (2006) Model for assessing applications concerning food fortification, Opinion of the
591 Panel on nutrition, dietetic products, novel food and allergy of the Norwegian
592 Scientific Committee for Food Safety, Norwegian Scientific Committee for Food
593 Safety, Oslo, Norway.
- 594 VKM. (2013a) Assessment and updating of the fortification model from 2006, Opinion of the
595 Panel on nutrition, dietetic products, novel food and allergy of the Norwegian
596 Scientific Committee for Food Safety Norwegian Scientific Committee for Food
597 Safety, Oslo, Norway. ISBN: 978-82-8259-069-3.
- 598 VKM. (2013b) Assessment of vitamin A and D in food supplements, Opinion of the Panel on
599 nutrition, dietetic products, novel food and allergy of the Norwegian Scientific
600 Committee for Food Safety, Norwegian Scientific Committee for Food Safety Oslo
601 Norway. ISBN: 978-82-8259-067-9.
- 602 VKM. (2014) Evaluation of tolerable upper intake levels for vitamin D in children and
603 adolescents, Statement of the Norwegian Scientific Committee for Food Safety,
604 Norwegian Scientific Committee for Food Safety, Oslo, Norway. ISBN: 978-82-8259-
605 140-9.
- 606 VKM. (2015a) Risk assessment of beta-carotene in food supplements, Opinion of the Panel
607 on Nutrition, Dietetic Products, Novel Food and Allergy of the Norwegian Scientific
608 Committee for Food Safety, Norwegian Scientific Committee for Food Safety, Oslo,
609 Norway. ISBN: 978-82-8259-155-3.
- 610 VKM. (2015b) Risk assessment of folic acid in food supplements, Opinion of the Panel on
611 Nutrition, Dietetic Products, Novel Food and Allergy of the Norwegian Scientific
612 Committee for Food Safety, Norwegian Scientific Committee for Food Safety, Oslo,
613 Norway. ISBN: 978-82-8259-154-6.
- 614 VKM. (2016a) Assessment of dietary intake of vitamin C and calcium in the Norwegian
615 population, Opinion of the Panel on Nutrition, Dietetic Products, Novel Food and
616 Allergy of the Norwegian Scientific Committee for Food Safety, Norwegian Scientific
617 Committee for Food Safety, Oslo, VKM. ISBN: 978-82-8259-206-2.
- 618 VKM. (2016b) Risk assessment of magnesium in food supplements, Opinion of the Panel on
619 Nutrition, Dietetic Products, Novel Food and Allergy of the Norwegian Scientific
620 Committee for Food Safety, Norwegian Scientific Committee for Food Safety, Oslo,
621 Norway. ISBN: 978-82-8259-205-5.
- 622 VKM. (2017a) Assessment of copper intake in relation to tolerable upper intake levels,
623 Opinion of the Panel on Nutrition, Dietetic Products, Novel Food and Allergy of the
624 Norwegian Scientific Committee for Food Safety, Norwegian Scientific Committee for
625 Food Safety, Oslo, Norway. ISBN: 978-82-8259-276-5.
- 626 VKM. (2017b) Assessment of dietary intake of phosphorus in relation to tolerable upper
627 intake levels, Opinion of the Panel on Nutrition, Dietetic Products, Novel Food and
628 Allergy of the Norwegian Scientific Committee for Food Safety, Norwegian Scientific
629 Committee for Food Safety, Oslo, Norway. ISBN: 978-82-8259-275-8.

- 630 VKM. (2017c) Assessment of intake of nicotinic acid and nicotinamide in relation to tolerable
631 upper intake levels, Opinion of the Panel on Nutrition, Dietetic Products, Novel Food
632 and Allergy of the Norwegian Scientific Committee for Food Safety, Norwegian
633 Scientific Committee for Food Safety, Oslo, Norway. ISBN: 978-82-8259-284-0.
- 634 VKM. (2017d) Assessment of iron intake in relation to tolerable upper intake levels, Opinion
635 of the Panel on Nutrition, Dietetic Products, Novel Food and Allergy of the Norwegian
636 Scientific Committee for Food Safety, Norwegian Scientific Committee for Food
637 Safety, Oslo Norway. ISBN: 978-82-8259-262-8.
- 638 VKM. (2017e) Assessment of selenium intake in relation to tolerable upper intake levels,
639 Opinion of the Panel on Nutrition, Dietetic Products, Novel Food and Allergy of the
640 Norwegian Scientific Committee for Food Safety, Norwegian Scientific Committee for
641 Food Safety, Oslo, Norway. ISBN: 978-82-8259-277-2.
- 642 VKM. (2017f) Assessment of vitamin B6 intake in relation to tolerable upper intake levels,
643 Opinion of the Panel on Nutrition, Dietetic Products, Novel Food and Allergy of the
644 Norwegian Scientific Committee for Food Safety, Norwegian Scientific Committee for
645 Food Safety, Oslo, Norway. ISBN: 978-82-8259-260-4.
- 646 VKM. (2017g) Assessment of vitamin E intake in relation to tolerable upper intake levels,
647 Opinion of the Panel on Nutrition, Dietetic Products, Novel Food and Allergy of the
648 Norwegian Scientific Committee for Food Safety, Norwegian Scientific Committee for
649 Food Safety, Oslo, Norway. ISBN: 978-82-8259-261-4.
- 650 VKM. (2017h) Assessment of zinc intake in relation to tolerable upper intake levels, Opinion
651 of the Panel on Nutrition, Dietetic Products, Novel Food and Allergy of the Norwegian
652 Scientific Committee for Food Safety, Norwegian Scientific Committee for Food Safety
653 Oslo, Norway. ISBN: 978-82-8259-263-5.
- 654 VKM. (2018a) Assessment of dietary intake of chromium (III) in relation to tolerable upper
655 intake level, Opinion of the Panel on Nutrition, Dietetic Products, Novel Food and
656 Allergy of the Norwegian Scientific Committee for Food and Environment, Norwegian
657 Scientific Committee for Food and Environment, Oslo, Norway. ISBN: 978-82-8259-
658 303-8.
- 659 VKM. (2018b) Assessment of dietary intake of manganese in relation to tolerable upper
660 intake level, Opinion of the Panel on Nutrition, Dietetic Products, Novel Food and
661 Allergy of the Norwegian Scientific Committee for Food and Environment, Norwegian
662 Scientific Committee on Food and Environment, Oslo, Norway. ISBN: 978-82-8259-
663 299-4.
- 664 VKM. (2018c) Assessment of dietary intake of vitamin K and maximum limits for vitamin K in
665 food supplements, Opinion of the Panel on Nutrition, Dietetic Products, Novel Food
666 and Allergy of the Norwegian Scientific Committee for Food and Environment,
667 Norwegian Scientific Committee for Food and Environment, Oslo, Norway. ISBN: 978-
668 82-8259-300-7.
- 669 VKM. (2020) Benefit and risk assessment of iodization of household salt and salt used in
670 bread and bakery products, Scientific opinion of the Panel on Nutrition, Dietetic
671 Products, Novel Food and Allergy of the Norwegian Scientific Committee for Food and

672 Environment, Norwegian Scientific Committee for Food and Environment, Oslo,
673 Norway. ISBN: 978-82-8259-343-4.
674

675 Appendix I

676 **Tolerable Upper Intake Levels (ULs), Safe Upper Levels (SULs) and Guidance Levels (GLs) used for**
 677 **this fortification model**

Nutrients	Unit	Men ≥ 18 years	Women ≥ 18 years	Adolescents 13 years	Children 9 years	Children 4 years	Children 2 years	Infants 1 year	Reference
Vitamin A	RE	3000	3000	2000	1500	1100	800	800	VKM, 2013b based on SCF, 2002
β-carotene*	µg	4000	4000	3000	2000	2000	1000	1000	EFSA, 2018 (no UL); VKM, 2015a
Vitamin D	µg	100	100	50	25	25	25	25	VKM, 2013b; VKM, 2014
Tocopherol	mg	300	300	220	160	120	100	100	EFSA, 2018; VKM, 2017g
Thiamine	mg	50	50	34	25	20	15	15	Rasmussen et al., 2006
Riboflavin	mg	43	43	29	22	16	12	12	Rasmussen et al., 2006; EVM, 2003
Niacin	mg	900	900	500	350	220	150	150	EFSA, 2018; VKM, 2017c
Vitamin B12	µg	2000	2000	1330	1000	730	530	530	Rasmussen et al., 2006; EVM, 2003
Vitamin B6	mg	25	25	20	10	7	5	5	EFSA, 2018; VKM, 2017f
Folic acid*	µg	1000	1000	600	400	300	200	200	EFSA, 2018; VKM, 2015b
Vitamin C	mg	1000	1000	1200	1200	650	400	400	EFSA, 2018 (no UL); VKM, 2016a
Calcium	mg	2500	2500	2500	2500	2500	2500	2500	EFSA, 2018 (no UL in children and adolescents); VKM, 2016a
Magnesium*	mg	250	250	250	250	250	85	85	EFSA, 2018 (no UL in young children); VKM, 2016b
Iron	mg	50	50	30	20	14	10	10	EFSA, 2018 (no UL); VKM, 2017d
Zinc	mg	40	40	23	23	12	7	7	VKM, 2017h
Copper	mg	5	5	4	3	2	1	1	EFSA, 2018; VKM, 2017a
Selenium	µg	300	300	200	130	90	60	60	EFSA, 2018; VKM, 2017e
Iodine	µg	600	600	450	300	250	200	200	EFSA, 2018; VKM, 2020
Phosphorus	mg	3000	3000	3000	3000	3000	3000	3000	EFSA, 2018 (no UL); VKM, 2017b

Nutrients	Unit	Men ≥ 18 years	Women ≥ 18 years	Adolescents 13 years	Children 9 years	Children 4 years	Children 2 years	Infants 1 year	Reference
Vitamin K	µg	1000	1000	670	500	370	270	270	EFSA, 2018 (no UL); VKM, 2018c
Chromium	µg	21000	21000	13020	6930	6930	3570	3570	EFSA, 2018 (no UL); VKM, 2018a
Manganese	mg	0	0	0	0	0	0	0	EFSA, 2018 (no UL); VKM, 2018b

678 *The ULs for beta-carotene, folic acid and magnesium are only applicable for the amount of nutrients added as fortifying agents or in food supplements, and
679 not for those that are naturally present in foods/drinks.

681 Appendix II

682 All calculations in the fortification model

683 Intakes in the 95th percentile in all included age groups

684 **Table Appendix II-1:** Intakes in the 95th percentile in men above 70 years from the Skedsmo
685 study. Energy intake in the 95th percentile is approximately 2900 kcal.

Skedsmo study, men 70 +	UL/GL	95P intake from food/ drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
Retinol, µg	3000	1786	484	730	101	126	168	252	503
β-carotene, µg	4000	0	150	3850	531	664	885	1328	2655
Vitamin D, µg	100	19.5	16.5	64.0	9	11	15	22	44
Tocopherol, mg	300	16.8	16.8	266	37	46	61	92	184
Thiamine, mg	50	2.4	0.8	47	6	8	11	16	32
Riboflavin, mg	43	2.9	0.9	39	5	7	9	13	27
Niacin, mg	900	34.0	10.5	856	118	148	197	295	590
Vitamin B12, µg	2000	17.8	51	1931	266	333	444	666	1332
Vitamin B6, mg	25	2.6	1.2	21	3	4	5	7	15
Folic acid, µg	1000	0	161	839	116	145	193	289	579
Vitamin C, mg	1000	215	40	745	103	128	171	257	514
Calcium, mg	2500	1563	20	917	126	158	211	316	632
Magnesium, mg	250	0	170	80	11	14	18	28	55
Iron, mg	50	16.4	8	25.6	4	4	6	9	18
Zinc, mg	40	16.9	9.2	13.9	2	2	3	5	10
Copper, mg	5	1.7	1.1	2.2	0.3	0.4	0.5	0.7	1
Selenium, µg	300	129	33	138	19	24	32	48	95
Iodine, µg	600	624	76	-100	-14	-17	-23	-34	-69
Phosphorus, mg	3000	2381	0	619	85	107	142	213	427

686

687 **Table Appendix II-2:** Intakes in the 95th percentile in women above 70 years from the Skedsmo
688 study. Energy intake in the 95th percentile is approximately 2500 kcal.

Skedsmo study, women 70 +	UL/GL	95P intake from food/ drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
Retinol, µg	3000	1303	404	1293	207	259	345	517	1034
β-carotene, µg	4000	0	150	3850	616	770	1027	1540	3080
Vitamin D, µg	100	16.6	11.8	72	11	14	19	29	57
Tocopherol, mg	300	15.8	12.2	272	44	54	73	109	218

Skedsmo study, women 70 +	UL/GL	95P intake from food/ drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
					Thiamine, mg	50	1.9	0.81	47
Riboflavin, mg	43	2.3	1.1	40	6	8	11	16	32
Niacin, mg	900	28.5	10.2	861	138	172	230	345	689
Vitamin B12, µg	2000	16.1	0.79	1983	317	397	529	793	1586
Vitamin B6, mg	25	2.2	1.2	22	3	4	6	9	17
Folic acid, µg	1000	0	98	902	144	180	241	361	722
Vitamin C, mg	1000	219	50.6	731	117	146	195	292	584
Calcium, mg	2500	1291	186	1023	164	205	273	409	819
Magnesium, mg	250	0	49.9	200	32	40	53	80	160
Iron, mg	50	13	4.6	32	5	6	9	13	26
Zinc, mg	40	13	6.2	20	3	4	5	8	16
Copper, mg	5	1	1	2.5	0.4	0.5	0.7	1.0	2
Selenium, µg	300	99	23.5	177	28	35	47	71	142
Iodine, µg	600	424	57	119	19	24	32	47	95
Phosphorus, mg	3000	2005	0	995	159	199	265	398	796

689

690 **Table Appendix II-3:** Intakes in the 95th percentile in men 18-70 years from Norkost 3 and
691 DANSDA 2011-2013 (for vitamin K1 and chromium). Energy intake in the 95th percentile is
692 approximately 4200 kcal in Norkost 3 and 3900 kcal in DANSDA.

Norkost 3, men 18-70 y (data for vitamin K1 and chromium is from DANSDA)	UL/GL	95P intake from food/ drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
					Retinol, µg	3000	1691	467	842
β-carotene, µg	4000	0	1443	2557	244	304	406	609	1218
Vitamin D, µg	100	17.2	13.6	69.2	7	8	11	16	33
Tocopherol, mg	300	23	15	262	25	31	42	62	125
Thiamine, mg	50	3.1	1.2	45.66	4	5	7	11	22
Riboflavin, mg	43	3.6	1.4	37.98	4	5	6	9	18
Niacin, mg	900	46	14.7	839.4	80	100	133	200	400
Vitamin B12, µg	2000	19.1	6	1974.93	188	235	313	470	940
Vitamin B6, mg	25	3.3	1.8	19.9	2	2	3	5	9
Folic acid, µg	1000	0	172	827.9	79	99	131	197	394
Vitamin C, mg	1000	258	135	607	58	72	96	145	289
Calcium, mg	2500	2032	169	299	28	36	47	71	142
Magnesium, mg	250	0	79	171	16	20	27	41	81
Iron, mg	50	21	9.8	19.2	2	2	3	5	9
Zinc, mg	40	22	10.4	7.6	1	1	1	2	4
Copper, mg	5	2.4	1.3	1.3	0.1	0.2	0.2	0.3	1
Selenium, µg	300	129	43	128	12	15	20	30	61
Iodine, µg	600	569	117	-86	-8	-10	-14	-20	-41
Phosphorus, mg	3000	3153	126	-279	-27	-33	-44	-66	-133

Norkost 3, men 18-70 y (data for vitamin K1 and chromium is from DANSDA)	UL/GL	95P intake from food/drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
Vitamin K1, µg	1000	187	8	805	83	103	138	206	413
Chromium, µg	21000	64	21	20915	2145	2681	3575	5363	10726

693

694 **Table Appendix II-4:** Intakes in the 95th percentile in women 18-70 years from Norkost 3 and
695 DANSDA 2011-2013 (for vitamin K1 and chromium). Energy intake in the 95th percentile is
696 approximately 2900 kcal in both Norkost 3 and DANSDA.

Norkost 3, women 18-70 y (data for vitamin K1 and chromium is from DANSDA)	UL/GL	95P intake from food/drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
Retinol, µg	3000	1285	386	1329	183	229	306	458	917
β-carotene, µg	4000	0	1556	2444	337	421	562	843	1686
Vitamin D, µg	100	12.3	11.4	76.3	11	13	18	26	53
Tocopherol, mg	300	18	12	270	37	47	62	93	186
Thiamine, mg	50	2.3	1.6	46.14	6	8	11	16	32
Riboflavin, mg	43	2.6	1.8	38.56	5	7	9	13	27
Niacin, mg	900	32	12.4	855.6	118	148	197	295	590
Vitamin B12, µg	2000	13.1	12	1974.9	272	341	454	681	1362
Vitamin B6, mg	25	2.5	2.1	20.4	3	4	5	7	14
Folic acid, µg	1000	0	149	851	117	147	196	293	587
Vitamin C, mg	1000	244	119	637	88	110	146	220	439
Calcium, mg	2500	1500	276	724	100	125	166	250	499
Magnesium, mg	250	0	111	139	19	24	32	48	96
Iron, mg	50	17	17	16	2	3	4	6	11
Zinc, mg	40	16	9	15	2	3	3	5	10
Copper, mg	5	1.9	1.1	2	0.3	0.3	0.5	1	1
Selenium, µg	300	94	36	170	23	29	39	59	117
Iodine, µg	600	421	95	84	12	14	19	29	58
Phosphorus, mg	3000	2282	78	640	88	110	147	221	441
Vitamin K1, µg	1000	179	8	813	112	140	187	280	561
Chromium, µg	21000	46	21	20933	2887	3609	4812	7218	14437

697

698 **Table Appendix II-5:** Intakes in the 95th percentile in 13-yea-olds from Ungkost 3 and DANSDA
699 2011-2013 (for vitamin K1 and chromium). Energy intake in the 95th percentile is approximately
700 3100 kcal in Ungkost 3 and 3200 kcal in DANSDA.

Ungkost 3, 13-year-olds (data for vitamin K1 and chromium is from DANSDA)	UL/GL	95P intake from food/drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
					Retinol, µg	2000	1486	187	327
β-carotene, µg	3000	0	0	3000	387	484	645	968	1935
Vitamin D, µg	50	10.7	8	31	4	5	7	10	20
Tocopherol, mg	220	16	8.2	196	25	32	42	63	126
Thiamine, mg	34	2.1	0.6	31	4	5	7	10	20
Riboflavin, mg	29	2.4	0.6	26	3	4	6	8	17
Niacin, mg	500	28	9.6	462	60	75	99	149	298
Vitamin B12, µg	1330	10	1.4	1319	170	213	284	425	851
Vitamin B6, mg	20	2.1	0.87	17	2	3	4	5	11
Folic acid, µg	600	0	102	498	64	80	107	161	321
Vitamin C, mg	1200	117	51	1032	133	166	222	333	666
Calcium, mg	2500	1592	375	533	69	86	115	172	344
Magnesium, mg	250	0	71	179	23	29	38	58	115
Iron, mg	30	14	7.4	9	1	1	2	3	6
Zinc, mg	23	17	3.9	2	0	0.3	0.5	1	1
Copper, mg	4	1.5	0.29	2	0	0.4	0.5	1	1
Selenium, µg	200	62	17	121	16	20	26	39	78
Iodine, µg	450	230	65	155	20	25	33	50	100
Phosphorus, mg	3000	2257	0	743	96	120	160	240	479
Vitamin K1, µg	670	111	6	553	69	86	115	173	346
Chromium, µg	13020	49	14	12957	1620	2025	2699	4049	8098

701

702 **Table Appendix II-6:** Intakes in the 95th percentile in 9-year-olds from Ungkost 3 and DANSDA
703 2011-2013 (for vitamin K1 and chromium). Energy intake in the 95th percentile is approximately
704 3100 kcal in Ungkost 3 and 3200 kcal in DANSDA.

Ungkost 3, 9-year-olds (data for vitamin K1 and chromium is from DANSDA)	UL/GL	95P intake from food/drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
					Retinol, µg	1500	1154	188	158
β-carotene, µg	2000	0	0	2000	320	400	533	800	1600
Vitamin D, µg	25	8.8	6.7	10	2	2	3	4	8
Tocopherol, mg	160	13.7	6.7	140	22	28	37	56	112
Thiamine, mg	25	1.7	0.6	23	4	5	6	9	18
Riboflavin, mg	22	2.2	0.7	19	3	4	5	8	15
Niacin, mg	350	22.3	7.1	321	51	64	85	128	256
Vitamin B12, µg	1000	9.4	0.9	990	158	198	264	396	792
Vitamin B6, mg	10	1.7	0.7	8	1	2	2	3	6
Folic acid, µg	400	0	68	332	53	66	89	133	266
Vitamin C, mg	1200	116	39	1045	167	209	279	418	836
Calcium, mg	2500	1389	0	1111	178	222	296	444	889

Ungkost 3, 9-year-olds (data for vitamin K1 and chromium is from DANSDA)	UL/GL	95P intake from food/drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
					Magnesium, mg	250	0	59	191
Iron, mg	20	12.3	6	2	0.3	0.3	0.5	1	1
Zinc, mg	23	16	4	3	0.5	1	1	1	2
Copper, mg	3	1.4	0.3	1	0.2	0.3	0.3	1	1
Selenium, µg	130	58	17	55	9	11	15	22	44
Iodine, µg	300	217	69	14	2	3	4	6	11
Phosphorus, mg	3000	1996	0	1004	161	201	268	402	803
Vitamin K1, µg	500	89	5	406	62	78	104	156	312
Chromium, µg	6930	37	8	6885	1059	1324	1765	2648	5296

705

706 **Table Appendix II-7:** Intakes in the 95th percentile in 4-year-olds from Ungkost 3 and DANSDA
707 2011-2013 (for vitamin K1 and chromium). Energy intake in the 95th percentile is approximately
708 2000 kcal in Ungkost 3 and 2600 kcal in DANSDA.

Ungkost 3, 4-year-olds (data for vitamin K1 and chromium is from DANSDA)	UL/GL	95P intake from food/drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
					Retinol, µg	1100	1079	193	-172
β-carotene, µg	2000	0	250	1750	350	438	583	875	1750
Vitamin D, µg	25	9.9	7.2	7.9	2	2	3	4	8
Tocopherol, mg	120	13.6	7	99.4	20	25	33	50	99
Thiamine, mg	20	1.4	0.6	18	4	5	6	9	18
Riboflavin, mg	16	1.9	0.7	13.4	3	3	4	7	13
Niacin, mg	220	20	7.7	192.3	38	48	64	96	192
Vitamin B12, µg	730	8.5	1.7	720	144	180	240	360	720
Vitamin B6, mg	7	1.9	0.8	4.3	1	1	1	2	4
Folic acid, µg	300	0	55	245	49	61	82	123	245
Vitamin C, mg	650	121	44	485	97	121	162	243	485
Calcium, mg	2500	1243	361	896	179	224	299	448	896
Magnesium, mg	250	0	136	114	23	29	38	57	114
Iron, mg	14	10.2	12	-8.2	-2	-2	-3	-4	-8
Zinc, mg	12	12.1	5	-5.1	-1	-1	-2	-3	-5
Copper, mg	2	1.2	1	-0.2	0.0	-0.1	-0.1	-0.1	-0.2
Selenium, µg	90	51	27	12	2	3	4	6	12
Iodine, µg	250	198	64	-12	-2	-3	-4	-6	-12
Phosphorus, mg	3000	1662	502	836	167	209	279	418	836
Vitamin K1, µg	370	89	5	276	42	53	71	106	212
Chromium, µg	6930	37	8	6885	1059	1324	1765	2648	5296

709

710 **Table Appendix II-8:** Intakes in the 95th percentile in 2-year-olds from Småbarnskost 3. Energy
 711 intake in the 95th percentile is approximately 2000 kcal.

Småbarnskost 3, 2-year-olds	UL/GL	95P intake from food/ drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
					Retinol, µg	800	1922	216	-1338
β-carotene, µg	1000	0	0	1000	200	250	333	500	1000
Vitamin D, µg	25	8.2	9.9	6.9	1	2	2	3	7
Tocopherol, mg	100	12.7	8.6	78.7	16	20	26	39	79
Thiamine, mg	15	1.7	0.6	12.7	3	3	4	6	13
Riboflavin, mg	12	2.4	0.7	8.9	2	2	3	4	9
Niacin, mg	150	14.6	8.1	127.3	25	32	42	64	127
Vitamin B12, µg	530	7.3	1.3	521	104	130	174	261	521
Vitamin B6, mg	5	1.7	0.7	2.6	1	1	1	1	3
Folic acid, µg	200	0	77	123	25	31	41	62	123
Vitamin C, mg	400	138	39	223	45	56	74	112	223
Calcium, mg	2500	1574	251	675	135	169	225	338	675
Magnesium, mg	85	0	60	25	5	6	8	13	25
Iron, mg	10	11.6	7.4	-9	-2	-2	-3	-5	-9
Zinc, mg	7	11.9	5.6	-10.5	-2	-3	-4	-5	-11
Copper, mg	1	1.1	0.4	-0.5	-0.1	-0.1	-0.2	-0.3	-1
Selenium, µg	60	53	24	-17	-3	-4	-6	-9	-17
Iodine, µg	200	253	96	-149	-30	-37	-50	-75	-149
Phosphorus, mg	3000	1935	0	1065	213	266	355	533	1065

712

713 **Table Appendix II-9:** Intakes in the 95th percentile in 1-year-olds from Spedkost 3 (non-breastfeds
 714 only). Energy intake in the 95th percentile is approximately 2000 kcal.

Spedkost 3 (non- breastfeds only), 1-year-olds	UL/GL	95P intake from food/ drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
					Retinol, µg	800	2394	215	-1809
β-carotene, µg	1000	0	0	1000	200	250	333	500	1000
Vitamin D, µg	25	16.8	9.6	-1.4	-0.3	-0.4	-0.5	-0.7	-1.4
Tocopherol, mg	100	20	7.9	72.1	14	18	24	36	72
Thiamine, mg	15	2.4	0.7	11.9	2	3	4	6	12
Riboflavin, mg	12	2.3	0.8	8.9	2	2	3	4	9
Niacin, mg	150	19.4	8.9	121.7	24	30	41	61	122
Vitamin B12, µg	530	6.1	1.2	523	105	131	174	261	523
Vitamin B6, mg	5	1.9	0.8	2.3	0.5	1	1	1	2
Folic acid, µg	200	0	89	111	22	28	37	56	111
Vitamin C, mg	400	217	39.8	143.2	29	36	48	72	143
Calcium, mg	2500	1358	250	892	178	223	297	446	892

Spedkost 3 (non-breastfeds only), 1-year-olds	UL/GL	95P intake from food/drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
Magnesium, mg	85	0	65.4	19.6	4	5	7	10	20
Iron, mg	10	22.4	13.6	-26	-5	-7	-9	-13	-26
Zinc, mg	7	13.2	6.1	-12.3	-2	-3	-4	-6	-12
Copper, mg	1	1.3	0.4	-0.7	-0.1	-0.2	-0.2	-0.4	-1
Selenium, µg	60	52	26	-18	-4	-5	-6	-9	-18
Iodine, µg	200	267	104	-171	-34	-43	-57	-86	-171
Phosphorus, mg	3000	1648	0	1352	270	338	451	676	1352

715

716 **Intakes in the 90th percentile in all included age groups**

717 **Table Appendix II-10:** Intakes in the 90th percentile in men above 70 years from the Skedsmo
718 study. Energy intake in the 90th percentile is approximately 2700 kcal.

Skedsmo study, men 70 +	UL/GL	90P intake from food/drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
Retinol, µg	3000	1190	484	1326	196	246	327	491	982
β-carotene, µg	4000	0	150	3850	570	713	951	1426	2852
Vitamin D, µg	100	15.9	16.5	68	10	13	17	25	50
Tocopherol, mg	300	15.8	16.8	267	40	50	66	99	198
Thiamine, mg	50	2.1	0.81	47	7	9	12	17	35
Riboflavin, mg	43	2.6	0.94	39	6	7	10	15	29
Niacin, mg	900	30	10.5	859	127	159	212	318	636
Vitamin B12, µg	2000	15	50.8	1934	286	358	477	716	1432
Vitamin B6, mg	25	2.3	1.2	21	3	4	5	8	16
Folic acid, µg	1000	0	161	839	124	155	207	311	622
Vitamin C, mg	1000	163	40	797	118	148	197	295	591
Calcium, mg	2500	1263	20	1217	180	225	301	451	902
Magnesium, mg	250	0	170	80	12	15	20	30	59
Iron, mg	50	15	8	27	4	5	7	10	20
Zinc, mg	40	14	9.2	16	2	3	4	6	12
Copper, mg	5	1.6	1.1	2.3	0.3	0.4	0.6	0.9	2
Selenium, µg	300	107	33	160	24	30	39	59	118
Iodine, µg	600	448	76	76	11	14	19	28	56
Phosphorus, mg	3000	2206	0	794	118	147	196	294	588

719

720 **Table Appendix II-11:** Intakes in the 90th percentile in women above 70 years from the Skedsmo
721 study. Energy intake in the 90th percentile is approximately 2300 kcal.

Skedsmo study, women 70 +	UL/GL	90P intake from food/ drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
					Retinol, µg	3000	1012	404	1584
β-carotene, µg	4000	0	150	3850	670	837	1116	1674	3348
Vitamin D, µg	100	13.9	11.8	74	13	16	22	32	65
Tocopherol, mg	300	14.6	12.2	273	48	59	79	119	238
Thiamine, mg	50	1.7	0.81	47	8	10	14	21	41
Riboflavin, mg	43	2.0	1.1	40	7	9	12	17	35
Niacin, mg	900	25.6	10.2	864	150	188	250	376	751
Vitamin B12, µg	2000	13	0.79	1987	345	432	576	864	1727
Vitamin B6, mg	25	1.9	1.2	22	4	5	6	10	19
Folic acid, µg	1000	0	98	902	157	196	261	392	784
Vitamin C, mg	1000	185	50.6	764	133	166	221	332	664
Calcium, mg	2500	1109	186	1205	210	262	349	524	1048
Magnesium, mg	250	0	49.9	200	35	44	58	87	174
Iron, mg	50	11.9	4.6	34	6	7	10	15	29
Zinc, mg	40	11.9	6.2	22	4	5	6	10	19
Copper, mg	5	1.3	1	2.7	0.5	0.6	0.8	1.2	2
Selenium, µg	300	82	23.5	195	34	42	56	85	169
Iodine, µg	600	350	57	193	33	42	56	84	167
Phosphorus, mg	3000	1802	0	1198	208	260	347	521	1042

722

723 **Table Appendix II-12:** Intakes in the 90th percentile in men 18-70 years from Norkost 3 and
724 DANSDA 2011-2013 (for vitamin K1 and chromium). Energy intake in the 90th percentile is
725 approximately 3700 kcal in Norkost 3 and 3600 kcal in DANSDA.

Norkost 3, men 18-70 y (data for vitamin K1 and chromium is from DANSDA)	UL/GL	90P intake from food/ drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
					Retinol, µg	3000	1353	467	1180
β-carotene, µg	4000	0	1443	2557	276	346	461	691	1382
Vitamin D, µg	100	12.9	13.6	73.5	8	10	13	20	40
Tocopherol, mg	300	20	15	265	29	36	48	72	143
Thiamine, mg	50	2.8	1.2	46.03	5	6	8	12	25
Riboflavin, mg	43	3.2	1.4	38.39	4	5	7	10	21
Niacin, mg	900	39.5	14.7	845.8	91	114	152	229	457
Vitamin B12, µg	2000	15.5	6	1978.53	214	267	356	535	1069
Vitamin B6, mg	25	2.9	1.8	20.3	2	3	4	5	11
Folic acid, µg	1000	0	172	827.9	90	112	149	224	448
Vitamin C, mg	1000	212	135	653	71	88	118	176	353
Calcium, mg	2500	1676	169	655	71	89	118	177	354
Magnesium, mg	250	0	79	171	18	23	31	46	92
Iron, mg	50	18	9.8	22.2	2	3	4	6	12

Norkost 3, men 18-70 y (data for vitamin K1 and chromium is from DANSDA)	UL/GL	90P intake from food/ drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
					Zinc, mg	40	20	10.4	9.6
Copper, mg	5	2.1	1.3	1.6	0.2	0.2	0.3	0.4	1
Selenium, µg	300	109	43	148	16	20	27	40	80
Iodine, µg	600	458	117	25	3	3	5	7	14
Phosphorus, mg	3000	2838	126	36	4	5	6	10	19
Vitamin K1, µg	1000	158	8	834	93	116	154	232	463
Chromium, µg	21000	57	21	20922	2325	2906	3874	5812	11623

726

727 **Table Appendix II-13:** Intakes in the 90th percentile in women 18-70 years from Norkost 3 and
728 DANSDA 2011-2013 (for vitamin K1 and chromium). Energy intake in the 90th percentile is
729 approximately 2600 kcal in Norkost 3 and 2700 kcal in DANSDA.

Norkost 3, women 18-70 y (data for vitamin K1 and chromium is from DANSDA)	UL/GL	90P intake from food/ drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
					Retinol, µg	3000	928	386	1686
β-carotene, µg	4000	0	1556	2444	376	470	627	940	1880
Vitamin D, µg	100	9.8	11.4	78.8	12	15	20	30	61
Tocopherol, mg	300	15	12	273	42	53	70	105	210
Thiamine, mg	50	2.0	1.6	46.42	7	9	12	18	36
Riboflavin, mg	43	2.4	1.8	38.8	6	7	10	15	30
Niacin, mg	900	28.5	12.4	859.1	132	165	220	330	661
Vitamin B12, µg	2000	10.8	12	1977.2	304	380	507	760	1521
Vitamin B6, mg	25	2.2	2.1	20.7	3	4	5	8	16
Folic acid, µg	1000	0	149	851	131	164	218	327	655
Vitamin C, mg	1000	206	119	675	104	130	173	260	519
Calcium, mg	2500	1275	276	949	146	183	243	365	730
Magnesium, mg	250	0	111	139	21	27	36	53	107
Iron, mg	50	15	17	18	3	3	5	7	14
Zinc, mg	40	14	9	17	3	3	4	7	13
Copper, mg	5	1.7	1.1	2.2	0.3	0.4	1	1	2
Selenium, µg	300	80	36	184	28	35	47	71	142
Iodine, µg	600	340	95	165	25	32	42	63	127
Phosphorus, mg	3000	2082	78	840	129	162	215	323	646
Vitamin K1, µg	1000	143	8	849	126	157	210	314	629
Chromium, µg	21000	42	21	20937	3102	3877	5170	7754	15509

730

731 **Table Appendix II-14:** Intakes in the 90th percentile in 13-year-olds from Ungkost 3 and DANSDA
 732 2011-2013 (for vitamin K1 and chromium). Energy intake in the 90th percentile is approximately
 733 2700 kcal in Ungkost 3 and 2900 kcal in DANSDA.

Ungkost 3, 13-year-olds (data for vitamin K1 and chromium is from DANSDA)	UL/GL	90P intake from food/drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
Retinol, µg	2000	945	187	868	129	161	214	321	643
β-carotene, µg	3000	0	0	3000	444	556	741	1111	2222
Vitamin D, µg	50	8.1	8	34	5	6	8	13	25
Tocopherol, mg	220	14	8.2	198	29	37	49	73	147
Thiamine, mg	34	1.8	0.6	32	5	6	8	12	23
Riboflavin, mg	29	2.1	0.6	26	4	5	6	10	19
Niacin, mg	500	24	9.6	466	69	86	115	173	345
Vitamin B12, µg	1330	8.5	1.4	1320	196	244	326	489	978
Vitamin B6, mg	20	1.8	0.87	17	3	3	4	6	13
Folic acid, µg	600	0	102	498	74	92	123	184	369
Vitamin C, mg	1200	94	51	1055	156	195	260	391	781
Calcium, mg	2500	1366	375	759	112	141	187	281	562
Magnesium, mg	250	0	71	179	27	33	44	66	133
Iron, mg	30	12	7.4	11	2	2	3	4	8
Zinc, mg	23	15	3.9	4	1	1	1	2	3
Copper, mg	4	1.4	0.29	2	0.3	0.4	1	1	2
Selenium, µg	200	55	17	128	19	24	32	47	95
Iodine, µg	450	194	65	191	28	35	47	71	141
Phosphorus, mg	3000	1995	0	1005	149	186	248	372	744
Vitamin K1, µg	670	92	6	572	79	99	131	197	394
Chromium, µg	13020	42	14	12964	1788	2235	2980	4470	8941

734

735 **Table Appendix II-15:** Intakes in the 90h percentile in 9-year-olds from Ungkost 3 and DANSDA
 736 2011-2013 (for vitamin K1 and chromium). Energy intake in the 90th percentile is approximately
 737 2300 kcal in Ungkost 3 and 2400 kcal in DANSDA.

Ungkost 3, 9-year-olds (data for vitamin K1 and chromium is from DANSDA)	UL/GL	90P intake from food/drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
Retinol, µg	1500	830	188	482	84	105	140	210	419
β-carotene, µg	2000	0	0	2000	348	435	580	870	1739
Vitamin D, µg	25	7.3	6.7	11	2	2	3	5	10
Tocopherol, mg	160	12.1	6.7	141	25	31	41	61	123
Thiamine, mg	25	1.6	0.6	23	4	5	7	10	20
Riboflavin, mg	22	2.0	0.7	19	3	4	6	8	17
Niacin, mg	350	19	7.1	324	56	70	94	141	282
Vitamin B12, µg	1000	7.6	0.9	992	172	216	287	431	862

Ungkost 3, 9-year-olds (data for vitamin K1 and chromium is from DANSDA)	UL/GL	90P intake from food/drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
					Vitamin B6, mg	10	1.5	0.7	8
Folic acid, µg	400	0	68	332	58	72	96	144	289
Vitamin C, mg	1200	93	39	1068	186	232	310	464	929
Calcium, mg	2500	1214	0	1286	224	280	373	559	1118
Magnesium, mg	250	0	59	191	33	42	55	83	166
Iron, mg	20	11.2	6	3	0.5	1	1	1	2
Zinc, mg	23	14	4	5	1	1	1	2	4
Copper, mg	3	1.2	0.3	2	0.3	0.3	0.4	1	1
Selenium, µg	130	50	17	63	11	14	18	27	55
Iodine, µg	300	180	69	51	9	11	15	22	44
Phosphorus, mg	3000	1780	0	1220	212	265	354	530	1061
Vitamin K1, µg	500	72	5	423	71	88	118	176	353
Chromium, µg	6930	35	8	6887	1148	1435	1913	2870	5739

738

739 **Table Appendix II-16:** Intakes in the 90th percentile in 4-year-olds from Ungkost 3 and DANSDA
740 2011-2013 (for vitamin K1 and chromium). Energy intake in the 90th percentile is approximately
741 1800 kcal in Ungkost 3 and 2400 kcal in DANSDA.

Ungkost 3, 4-year-olds (data for vitamin K1 and chromium is from DANSDA)	UL/GL	90P intake from food/drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
					Retinol, µg	1100	848	193	59
β-carotene, µg	2000	0	250	1750	389	486	648	972	1944
Vitamin D, µg	25	8	7.2	9.8	2	3	4	5	11
Tocopherol, mg	120	12.6	7	100.4	22	28	37	56	112
Thiamine, mg	20	1.3	0.6	18.1	4	5	7	10	20
Riboflavin, mg	16	1.7	0.7	13.6	3	4	5	8	15
Niacin, mg	220	17	7.7	195.3	43	54	72	109	217
Vitamin B12, µg	730	6.9	1.7	721	160	200	267	401	802
Vitamin B6, mg	7	1.7	0.8	4.5	1	1	2	3	5
Folic acid, µg	300	0	55	245	54	68	91	136	272
Vitamin C, mg	650	104	44	502	112	139	186	279	558
Calcium, mg	2500	1109	361	1030	229	286	381	572	1144
Magnesium, mg	250	0	136	114	25	32	42	63	127
Iron, mg	14	9	12	-7	-2	-2	-3	-4	-8
Zinc, mg	12	10.4	5	-3.4	-1	-1	-1	-2	-4
Copper, mg	2	1.1	1	-0.1	0.0	0.0	0.0	-0.1	-0.1
Selenium, µg	90	45	27	18	4	5	7	10	20
Iodine, µg	250	160	64	26	6	7	10	14	29
Phosphorus, mg	3000	1510	502	988	220	274	366	549	1098
Vitamin K1, µg	370	72	5	293	49	61	81	122	244

Ungkost 3, 4-year-olds (data for vitamin K1 and chromium is from DANSDA)	UL/GL	90P intake from food/drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
					Chromium, µg	6930	35	8	6887

742

743 **Table Appendix II-17:** Intakes in the 90th percentile in 2-year-olds from Småbarnskost 3. Energy
744 intake in the 90th percentile is approximately 1800 kcal.

Småbarnskost 3, 2-year-olds	UL/GL	90P intake from food/drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
					Retinol, µg	800	1544	216	-960
β-carotene, µg	1000	0	0	1000	222	278	370	556	1111
Vitamin D, µg	25	7.2	9.9	7.9	2	2	3	4	9
Tocopherol, mg	100	11	8.6	80.4	18	22	30	45	89
Thiamine, mg	15	1.5	0.6	12.9	3	4	5	7	14
Riboflavin, mg	12	2.1	0.7	9.2	2	3	3	5	10
Niacin, mg	150	12.9	8.1	129	29	36	48	72	143
Vitamin B12, µg	530	6.4	1.3	522	116	145	193	290	580
Vitamin B6, mg	5	1.5	0.7	2.8	1	1	1	2	3
Folic acid, µg	200	0	77	123	27	34	46	68	137
Vitamin C, mg	400	114	39	247	55	69	91	137	274
Calcium, mg	2500	1363	251	886	197	246	328	492	984
Magnesium, mg	85	0	60	25	6	7	9	14	28
Iron, mg	10	10	7.4	-7.4	-2	-2	-3	-4	-8
Zinc, mg	7	10.6	5.6	-9.2	-2	-3	-3	-5	-10
Copper, mg	1	1	0.4	-0.4	-0.1	-0.1	-0.1	-0.2	-0.4
Selenium, µg	60	48	24	-12	-3	-3	-4	-7	-13
Iodine, µg	200	253	96	-149	-33	-41	-55	-83	-166
Phosphorus, mg	3000	1708	0	1292	287	359	479	718	1436

745

746 **Table Appendix II-18:** Intakes in the 90th percentile in 1-year-olds from Spedkost 3 (non-
747 breastfeds only). Energy intake in the 90th percentile is approximately 1700 kcal.

Spedkost 3 (non-breastfeds only), 1-year-olds	UL/GL	90P intake from food/drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
					Retinol, µg	800	1852	215	-1267
β-carotene, µg	1000	0	0	1000	235	294	392	588	1176
Vitamin D, µg	25	15.3	9.6	0.1	0.0	0.0	0.0	0.1	0.1

Spedkost 3 (non-breastfeds only), 1-year-olds	UL/GL	90P intake from food/drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
Tocopherol, mg	100	17.5	7.9	74.6	18	22	29	44	88
Thiamine, mg	15	2.1	0.7	12.2	3	4	5	7	14
Riboflavin, mg	12	2	0.8	9.2	2	3	4	5	11
Niacin, mg	150	17.3	8.9	123.8	29	36	49	73	146
Vitamin B12, µg	530	5.2	1.2	524	123	154	205	308	616
Vitamin B6, mg	5	1.7	0.8	2.5	1	1	1	1	3
Folic acid, µg	200	0	89	111	26	33	44	65	131
Vitamin C, mg	400	194	39.8	166.2	39	49	65	98	196
Calcium, mg	2500	1166	250	1084	255	319	425	638	1275
Magnesium, mg	85	0	65.4	19.6	5	6	8	12	23
Iron, mg	10	19.6	13.6	-23.2	-5	-7	-9	-14	-27
Zinc, mg	7	11.6	6.1	-10.7	-3	-3	-4	-6	-13
Copper, mg	1	1.2	0.4	-0.6	-0.1	-0.2	-0.2	-0.4	-1
Selenium, µg	60	47	26	-13	-3	-4	-5	-8	-15
Iodine, µg	200	215	104	-119	-28	-35	-47	-70	-140
Phosphorus, mg	3000	1454	0	1546	364	455	606	909	1819

748

749 **Intakes in the 75th percentile in all included age groups**

750 **Table Appendix II-19:** Intakes in the 75th percentile in men above 70 years from the Skedsmo
751 study. Energy intake in the 75th percentile is approximately 2300 kcal.

Skedsmo study, men 70 +	UL/GL	75P intake from food/drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
Retinol, µg	3000	837	484	1679	292	365	487	730	1460
β-carotene, µg	4000	0	150	3850	670	837	1116	1674	3348
Vitamin D, µg	100	9.3	16.5	74	13	16	21	32	64
Tocopherol, mg	300	12.2	16.8	271	47	59	79	118	236
Thiamine, mg	50	1.7	0.8	48	8	10	14	21	41
Riboflavin, mg	43	2.0	0.9	40	7	9	12	17	35
Niacin, mg	900	24.8	10.5	865	150	188	251	376	752
Vitamin B12, µg	2000	9	51	1940	337	422	562	844	1687
Vitamin B6, mg	25	1.8	1.2	22	4	5	6	10	19
Folic acid, µg	1000	0	161	839	146	182	243	365	730
Vitamin C, mg	1000	123	40	837	146	182	243	364	728
Calcium, mg	2500	971	20	1509	262	328	437	656	1312
Magnesium, mg	250	0	170	80	14	17	23	35	70
Iron, mg	50	11.6	8	30	5	7	9	13	26
Zinc, mg	40	11.7	9.2	19	3	4	6	8	17
Copper, mg	5	1.3	1.1	2.6	0.5	0.6	0.8	1	2

Skedsmo study, men 70 +	UL/GL	75P intake from food/drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
Selenium, µg	300	76	33	191	33	41	55	83	166
Iodine, µg	600	237	76	287	50	62	83	125	250
Phosphorus, mg	3000	1793	0	1207	210	262	350	525	1049

752

753 **Table Appendix II-20:** Intakes in the 75th percentile in women above 70 years from the Skedsmo
754 study. Energy intake in the 75th percentile is approximately 1900 kcal.

Skedsmo study, women 70 +	UL/GL	75P intake from food/drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
Retinol, µg	3000	724	404	1872	394	493	657	985	1971
β-carotene, µg	4000	0	150	3850	811	1013	1351	2026	4053
Vitamin D, µg	100	8.6	11.8	80	17	21	28	42	84
Tocopherol, mg	300	11.3	12.2	276	58	73	97	146	291
Thiamine, mg	50	1.4	0.81	48	10	13	17	25	50
Riboflavin, mg	43	1.7	1.1	40	8	11	14	21	42
Niacin, mg	900	19.9	10.2	870	183	229	305	458	916
Vitamin B12, µg	2000	8.3	0.79	1991	419	524	699	1048	2096
Vitamin B6, mg	25	1.6	1.2	22	5	6	8	12	23
Folic acid, µg	1000	0	98	902	190	237	316	475	949
Vitamin C, mg	1000	125	50.6	825	174	217	289	434	868
Calcium, mg	2500	851	186	1463	308	385	513	770	1540
Magnesium, mg	250	0	49.9	200	42	53	70	105	211
Iron, mg	50	10.0	4.6	35	7	9	12	19	37
Zinc, mg	40	9.9	6.2	24	5	6	8	13	25
Copper, mg	5	1.1	1	2.9	0.6	0.8	1.0	2	3
Selenium, µg	300	64	23.5	213	45	56	75	112	224
Iodine, µg	600	223	57	320	67	84	112	168	336
Phosphorus, mg	3000	1554	0	1446	304	380	507	761	1522

755

756 **Table Appendix II-21:** Intakes in the 75th percentile in men 18-70 years from Norkost 3 and
757 DANSDA 2011-2013 (for vitamin K1 and chromium). Energy intake in the 75th percentile is
758 approximately 3100 kcal in both Norkost 3 and DANSDA.

Norkost 3, men 18-70 y (data for vitamin K1 and chromium is from DANSDA)	UL/GL	75P intake from food/drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
Retinol, µg	3000	948	467	1585	205	256	341	511	1023

Norkost 3, men 18-70 y (data for vitamin K1 and chromium is from DANSDA)	UL/GL	75P intake from food/ drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
					β-carotene, µg	4000	0	1443	2557
Vitamin D, µg	100	8.4	13.6	78	10	13	17	25	50
Tocopherol, mg	300	15	15	270	35	44	58	87	174
Thiamine, mg	50	2.3	1.2	46.54	6	8	10	15	30
Riboflavin, mg	43	2.6	1.4	39.01	5	6	8	13	25
Niacin, mg	900	31.3	14.7	854	110	138	184	275	551
Vitamin B12, µg	2000	11	6	1983.3	256	320	427	640	1280
Vitamin B6, mg	25	2.3	1.8	20.9	3	3	4	7	13
Folic acid, µg	1000	0	172	827.9	107	134	178	267	534
Vitamin C, mg	1000	141	135	724	93	117	156	234	467
Calcium, mg	2500	1295	169	1036	134	167	223	334	668
Magnesium, mg	250	0	79	171	22	28	37	55	110
Iron, mg	50	15	9.8	25.2	3	4	5	8	16
Zinc, mg	40	16	10.4	13.6	2	2	3	4	9
Copper, mg	5	1.7	1.3	2	0.3	0.3	0.4	1	1
Selenium, µg	300	79	43	178	23	29	38	57	115
Iodine, µg	600	260	117	223	29	36	48	72	144
Phosphorus, mg	3000	2350	126	524	68	85	113	169	338
Vitamin K1, µg	1000	115	8	877	113	141	189	283	566
Chromium, µg	21000	49	21	20930	2701	3376	4501	6752	13503

759

760 **Table Appendix II-22:** Intakes in the 75th percentile in women 18-70 years from Norkost 3 and
761 DANSDA 2011-2013 (for vitamin K1 and chromium). Energy intake in the 75th percentile is
762 approximately 2300 kcal in both Norkost 3 and DANSDA.

Norkost 3, women 18-70 y (data for vitamin K1 and chromium is from DANSDA)	UL/GL	75P intake from food/ drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
					Retinol, µg	3000	653	386	1961
β-carotene, µg	4000	0	1556	2444	425	531	708	1063	2125
Vitamin D, µg	100	6.3	11.4	82.3	14	18	24	36	72
Tocopherol, mg	300	12	12	276	48	60	80	120	240
Thiamine, mg	50	1.7	1.6	46.75	8	10	14	20	41
Riboflavin, mg	43	1.9	1.8	39.3	7	9	11	17	34
Niacin, mg	900	22.9	12.4	864.7	150	188	251	376	752
Vitamin B12, µg	2000	7.8	12	1980.2	344	430	574	861	1722
Vitamin B6, mg	25	1.8	2.1	21.1	4	5	6	9	18
Folic acid, µg	1000	0	149	851	148	185	247	370	740
Vitamin C, mg	1000	147	119	734	128	160	213	319	638
Calcium, mg	2500	1027	276	1197	208	260	347	520	1041
Magnesium, mg	250	0	111	139	24	30	40	60	121

Norkost 3, women 18-70 y (data for vitamin K1 and chromium is from DANSDA)	UL/GL	75P intake from food/ drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
					Iron, mg	50	11.7	17	21.3
Zinc, mg	40	11.7	9	19.3	3	4	6	8	17
Copper, mg	5	1.4	1.1	2.5	0.4	1	1	1	2
Selenium, µg	300	60	36	204	35	44	59	89	177
Iodine, µg	600	189	95	316	55	69	92	137	275
Phosphorus, mg	3000	1781	78	1141	198	248	331	496	992
Vitamin K1, µg	1000	105	8	887	154	193	257	386	771
Chromium, µg	21000	36	21	20943	3642	4553	6070	9106	18211

763

764 **Table Appendix II-23:** Intakes in the 75th percentile in 13-year-olds from Ungkost 3 and DANSDA
765 2011-2013 (for vitamin K1 and chromium). Energy intake in the 75th percentile is approximately
766 2300 kcal in Ungkost 3 and 2400 kcal in DANSDA.

Ungkost 3, 13- year-olds (data for vitamin K1 and chromium is from DANSDA)	UL/GL	75P intake from food/ drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
					Retinol, µg	2000	506	187	1307
β-carotene, µg	3000	0	0	3000	522	652	870	1304	2609
Vitamin D, µg	50	4.9	8	37	6	8	11	16	32
Tocopherol, mg	220	11	8.2	201	35	44	58	87	175
Thiamine, mg	34	1.5	0.6	32	6	7	9	14	28
Riboflavin, mg	29	1.7	0.6	27	5	6	8	12	23
Niacin, mg	500	19	9.6	471	82	102	137	205	410
Vitamin B12, µg	1330	6.5	1.4	1322	230	287	383	575	1150
Vitamin B6, mg	20	1.4	0.87	18	3	4	5	8	15
Folic acid, µg	600	0	102	498	87	108	144	217	433
Vitamin C, mg	1200	61	51	1088	189	237	315	473	946
Calcium, mg	2500	1049	375	1076	187	234	312	468	936
Magnesium, mg	250	0	71	179	31	39	52	78	156
Iron, mg	30	10	7.4	13	2	3	4	5	11
Zinc, mg	23	13	3.9	6	1	1	2	3	5
Copper, mg	4	1.1	0.29	3	0.5	1	1	1	2
Selenium, µg	200	42	17	141	25	31	41	61	123
Iodine, µg	450	131	65	254	44	55	74	110	221
Phosphorus, mg	3000	1658	0	1342	233	292	389	583	1167
Vitamin K1, µg	670	69	6	595	99	124	165	248	496
Chromium, µg	13020	33	14	12973	2162	2703	3604	5405	10811

767

768 **Table Appendix II-24:** Intakes in the 75th percentile in 9-year-olds from Ungkost 3 and DANSDA
 769 2011-2013 (for vitamin K1 and chromium). Energy intake in the 75th percentile is approximately
 770 2000 kcal in Ungkost 3 and 2200 kcal in DANSDA.

Ungkost 3, 9-year-olds (data for vitamin K1 and chromium is from DANSDA)	UL/GL	75P intake from food/drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
Retinol, µg	1500	524	188	788	158	197	263	394	788
β-carotene, µg	2000	0	0	2000	400	500	667	1000	2000
Vitamin D, µg	25	4.8	6.7	14	3	3	5	7	14
Tocopherol, mg	160	10	6.7	143	29	36	48	72	143
Thiamine, mg	25	1.4	0.6	23	5	6	8	12	23
Riboflavin, mg	22	1.6	0.7	20	4	5	7	10	20
Niacin, mg	350	15.1	7.1	328	66	82	109	164	328
Vitamin B12, µg	1000	6	0.9	993	199	248	331	497	993
Vitamin B6, mg	10	1.2	0.7	8	2	2	3	4	8
Folic acid, µg	400	0	68	332	66	83	111	166	332
Vitamin C, mg	1200	60	39	1101	220	275	367	551	1101
Calcium, mg	2500	995	0	1505	301	376	502	753	1505
Magnesium, mg	250	0	59	191	38	48	64	96	191
Iron, mg	20	9.3	6	5	1	1	2	2	5
Zinc, mg	23	11	4	8	2	2	3	4	8
Copper, mg	3	1	0.3	2	0.3	0.4	1	1	2
Selenium, µg	130	40	17	73	15	18	24	37	73
Iodine, µg	300	131	69	100	20	25	33	50	100
Phosphorus, mg	3000	1547	0	1453	291	363	484	727	1453
Vitamin K1, µg	500	54	5	441	80	100	134	200	401
Chromium, µg	6930	30	8	6892	1253	1566	2088	3133	6265

771

772 **Table Appendix II-25:** Intakes in the 75th percentile in 4-year-olds from Ungkost 3 and DANSDA
 773 2011-2013 (for vitamin K1 and chromium). Energy intake in the 75th percentile is approximately
 774 1600 kcal in Ungkost 3 and 2200 kcal in DANSDA.

Ungkost 3, 4-year-olds (data for vitamin K1 and chromium is from DANSDA)	UL/GL	75P intake from food/drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
Retinol, µg	1100	596	193	311	78	97	130	194	389
β-carotene, µg	2000	0	250	1750	438	547	729	1094	2188
Vitamin D, µg	25	5.9	7.2	11.9	3	4	5	7	15
Tocopherol, mg	120	10.3	7	102.7	26	32	43	64	128
Thiamine, mg	20	1.1	0.6	18.3	5	6	8	11	23
Riboflavin, mg	16	1.5	0.7	13.8	3	4	6	9	17
Niacin, mg	220	14	7.7	198.3	50	62	83	124	248
Vitamin B12, µg	730	5.8	1.7	723	181	226	301	452	903

Ungkost 3, 4-year-olds (data for vitamin K1 and chromium is from DANSDA)	UL/GL	75P intake from food/drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
					Vitamin B6, mg	7	1.3		5.7
Folic acid, µg	300	0	55	245	61	77	102	153	306
Vitamin C, mg	650	76	44	530	133	166	221	331	663
Calcium, mg	2500	865	361	1274	319	398	531	796	1593
Magnesium, mg	250	0	136	114	29	36	48	71	143
Iron, mg	14	7.8	12	-5.8	-1	-2	-2	-4	-7
Zinc, mg	12	8.9	5	-1.9	-0.5	-1	-1	-1	-2
Copper, mg	2	0.9	1	0.1	0.0	0.0	0.0	0.1	0.1
Selenium, µg	90	38	27	25	6	8	10	16	31
Iodine, µg	250	131	64	55	14	17	23	34	69
Phosphorus, mg	3000	1313	502	1185	296	370	494	741	1481
Vitamin K1, µg	370	54	5	311	57	71	94	141	283
Chromium, µg	6930	30	8	6892	1253	1566	2088	3133	6265

775

776 **Table Appendix II-26:** Intakes in the 75th percentile in 2-year-olds from Småbarnskost 3. Energy
777 intake in the 75th percentile is approximately 1500 kcal.

Småbarnskost 3, 2-year-olds	UL/GL	75P intake from food/drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
					Retinol, µg	800	1063	216	-479
β-carotene, µg	1000	0	0	1000	267	333	444	667	1333
Vitamin D, µg	25	5.6	9.9	9.5	3	3	4	6	13
Tocopherol, mg	100	8.9	8.6	82.5	22	28	37	55	110
Thiamine, mg	15	1.2	0.6	13.2	4	4	6	9	18
Riboflavin, mg	12	1.7	0.7	9.6	3	3	4	6	13
Niacin, mg	150	10.8	8.1	131.1	35	44	58	87	175
Vitamin B12, µg	530	5.1	1.3	524	140	175	233	349	698
Vitamin B6, mg	5	1.2	0.7	3.1	1	1	1	2	4
Folic acid, µg	200	0	77	123	33	41	55	82	164
Vitamin C, mg	400	84	39	277	74	92	123	185	369
Calcium, mg	2500	1090	251	1159	309	386	515	773	1545
Magnesium, mg	85	0	60	25	7	8	11	17	33
Iron, mg	10	8.2	7.4	-5.6	-1	-2	-2	-4	-7
Zinc, mg	7	8.8	5.6	-7.4	-2	-2	-3	-5	-10
Copper, mg	1	0.8	0.4	-0.2	-0.1	-0.1	-0.1	-0.1	-0.3
Selenium, µg	60	39	24	-3	-1	-1	-1	-2	-4
Iodine, µg	200	253	96	-149	-40	-50	-66	-99	-199
Phosphorus, mg	3000	1417	0	1583	422	528	704	1055	2111

778

779 **Table Appendix II-27:** Intakes in the 75th percentile in 1-year-olds from Spedkost 3 (non-
 780 breastfeds only). Energy intake in the 75th percentile is approximately 1400 kcal.

Spedkost 3 (non-breastfeds only), 1-year-olds	UL/GL	75P intake from food/drink	Intake from suppl. (mean, users only)	Maximal amount (MA)	Max addition pr. 100 kcal (FA) E% from fortified foods				
					25	20	15	10	5
					Retinol, µg	800	1410	215	-825
β-carotene, µg	1000	0	0	1000	286	357	476	714	1429
Vitamin D, µg	25	12.0	9.6	3.4	1	1	2	2	5
Tocopherol, mg	100	14.3	7.9	77.8	22	28	37	56	111
Thiamine, mg	15	1.8	0.7	12.5	4	4	6	9	18
Riboflavin, mg	12	1.7	0.8	9.5	3	3	5	7	14
Niacin, mg	150	14.5	8.9	126.6	36	45	60	90	181
Vitamin B12, µg	530	4.3	1.2	525	150	187	250	375	749
Vitamin B6, mg	5	1.4	0.8	2.8	1	1	1	2	4
Folic acid, µg	200	0	89	111	32	40	53	79	159
Vitamin C, mg	400	156	39.8	204.2	58	73	97	146	292
Calcium, mg	2500	962	250	1288	368	460	613	920	1840
Magnesium, mg	85	0	65.4	19.6	6	7	9	14	28
Iron, mg	10	15.8	13.6	-19.4	-6	-7	-9	-14	-28
Zinc, mg	7	9.8	6.1	-8.9	-3	-3	-4	-6	-13
Copper, mg	1	1	0.4	-0.4	-0.1	-0.1	-0.2	-0.3	-1
Selenium, µg	60	39	26	-5	-1	-2	-2	-4	-7
Iodine, µg	200	164	104	-68	-19	-24	-32	-49	-97
Phosphorus, mg	3000	1211	0	1789	511	639	852	1278	2556

781